

Europäische
Patentamt

European Patent
Office

PCT/EP 03/1179
Office européen
des brevets

10/532740

REC'D 10 DEC 2003

WIPO

PCT



Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten internationalen Patentanmeldung überein.

The attached documents are exact copies of the international patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet international spécifiée à la page suivante.

Den Haag, den
The Hague,
La Haye, le

03. 12. 03

Der Präsident des Europäischen Patentamts
Im Auftrag
For the President of the European Patent Office
Le Président de l'Office européen des brevets
p.o.

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)


NATHALIE KUIPER

Patentanmeldung Nr.
Patent application no.
Demande de brevet n°

PCT/EP 02/12273 ✓

BEST AVAILABLE COPY

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

19

Anmeldung Nr.:
Application no.:
Demande n°:

PCT/EP 02/12273 ✓

Anmelder:
Applicant(s):
Demandeur(s):

1. JANSSEN PHARMACEUTICA N.V. - Beerse, Belgium
2. PEETERS, Pieter, Johan - Beerse, Belgium (US only)
3. GÖHLMANN, Hinrich, Wilhelm, Helmut - Beerse, Belgium (US only)

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:

STRESS RELATED GENES

Anmeldetag:
Date of filing:
Date de dépôt:

31 October 2002 (31.10.2002) ✓

In Anspruch genommene Priorität(en)
Priority(ies) claimed
Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Benennung von Vertragsstaaten : Siehe Formblatt PCT/RO/101 (beigefügt)
Designation of contracting states : See Form PCT/RO/101 (enclosed)
Désignation d'états contractants : Voir Formulaire PCT/RO/101 (ci-joint)

Bemerkungen:
Remarks:
Remarques:

Further applicants

4. SWAGEMAKERS, Sigrid, Maria, Alice - Beerse, Belgium (US only)

PCT REQUEST

Original (for SUBMISSION) - printed on 31.10.2002 03:55:30 PM

III-2	Applicant and/or inventor	applicant and inventor
III-2-1	This person is:	US only
III-2-2	Applicant for	GÖHLMANN, Hinrich, Wilhelm, Helmut
III-2-4	Name (LAST, First)	Janssen Pharmaceutica N.V.
III-2-5	Address:	B-2340 Beerse Belgium
III-2-6	State of nationality	DE
III-2-7	State of residence	NL
III-3	Applicant and/or inventor	applicant and inventor
III-3-1	This person is:	US only
III-3-2	Applicant for	SWAGENAKERS, Sigrid, Maria, Alice
III-3-4	Name (LAST, First)	Janssen Pharmaceutica N.V.
III-3-5	Address:	B-2340 Beerse Belgium
III-3-6	State of nationality	NL
III-3-7	State of residence	NL
IV-1	Agent or common representative; or address for correspondence The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:	common representative
IV-1-1	Name	JANSSEN PHARMACEUTICA N.V.
IV-1-2	Address:	Turnhoutseweg 30 B-2340 Beerse Belgium
IV-1-3	Telephone No.	32+14 60 21 86
IV-1-4	Facsimile No.	32+14 60 54 91
IV-1-5	e-mail	patents@janbe.jnj.com
V	Designation of States	
V-1	Regional Patent (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	EP: AT BE BG CH LI CY CZ DE DK EE ES FI FR GB GR IE IT LU MC NL PT SE SK TR and any other State which is a Contracting State of the European Patent Convention and of the PCT
V-2	National Patent (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	US

STRESS RELATED GENES

5 The present invention relates generally to therapy and diagnosis of depression. In particular this invention relates to the polypeptides as well as to the polynucleotides encoding these polypeptides, wherein said polypeptides are shown to play a central role in mediating the cellular response to corticotropin releasing hormone. These polypeptides and polynucleotides are useful in the diagnosis, treatment and/or prevention of depression.

10

BACKGROUND OF THE INVENTION

15 Recent socioeconomic analyses found that depression is a leading cause of disability and a major risk factor for development of other diseases. Moreover, on a world-wide scale depression is underdiagnosed and undertreated. Current antidepressant drugs have proven to be effective, but are burdened with slow onset of action and side effects. Above this, it is still unclear by which pharmacological mode of action they exert their clinical effects. Hypothesis-driven research based upon the corticosteroid receptor hypothesis of depression has led to a novel concept focusing on brain neuropeptide receptors, specifically the corticotropin-releasing hormone (CRH) receptor as drug target.

20 Corticotropin releasing hormone (CRH), a 41-amino acid polypeptide plays a central role in the regulation of the hypothalamic-pituitary-adrenal axis, mediating the endocrine responses to various stressors. Hypothalamic neurons release CRH into the hypophyseal portal system in response to stress, stimulating the secretion and biosynthesis of pituitary adrenocorticotropin (ACTH) leading to increased adrenal glucocorticoid production (1). Several clinical and preclinical studies point towards a causal role for alterations in the CRH system in the development of depression (2). The first studies with CRH in humans showed that the ACTH response to CRH is blunted in depressed patients, reflecting a CRH receptor desensitization secondary to continuously increased hypothalamic CRH secretion (3;4). In support of blunted ACTH response as consequence of increased CRH release is the finding of elevated CRH levels in cerebrospinal fluid of patients with depression. Other findings strengthening this notion of CRH hypersecretion in the depressed state are an increased number of CRH secreting

25
30
35

neurons and a decreased number of CRH receptors in suicide victims who suffered from depression (5;6).

For CRH two high affinity receptors have been described, CRH-R1 and CRH-R2, both of which exist in several splice variant forms. Activation of these receptors by CRH results in G_s-mediated stimulation of adenylyl cyclase leading to increased levels of intracellular cAMP. This in itself will activate cAMP dependent protein kinase A (PKA) and ultimately result in increased cytosolic levels of cAMP and Ca²⁺. The increased levels of cAMP and Ca²⁺ lead to the activation of several other additional kinases such as Ca²⁺/calmodulin-dependent kinase II (CAMKII) and p42/p44 mitogen activated kinases (MAPK). As a result the Ca²⁺/cAMP response element binding protein (CREB) is phosphorylated and this in turn will regulate the transcription of genes containing cAMP response elements (CRE) in their promoter region. Examples of such genes shown to be involved in the modulation of CRH signaling include *c-fos*, macrophage migration-inhibitory factor gene *Mif*, orphan nuclear receptors *Nurr77* and *Nurr1*.

Notwithstanding the fact that the downstream pathways for CRH activated receptors were extensively studied and led to the identification of a number of genes involved in the signaling cascade, a major area is unexplored. It was thus an object of the present invention to explore the transcriptional response to CRH stimulation at a genome wide level in order to identify further genes involved in the corticotropin-releasing hormone receptor activated gene network. The polypeptides thus identified and the polynucleotides encoding said polypeptides provide new chances for drug development as drug targets through screening techniques, or are useful in the diagnosis, prevention and/or treatment of depression.

BRIEF DESCRIPTION OF THE DRAWING

Table 1: List of proteins which modulate CRH signaling.

Figure 1: *c-fos* mRNA levels as assessed by quantitative RT-PCR normalized against β -actin mRNA levels (taken as 100%) in AtT-20 cells treated with DMSO, CRH, CRH+R121919 or R121919 for different time points (hours).

Figure 2: Correspondence analysis applied on normalized microarray data for all time points and treatments. Squares depict different samples whereas circles depict genes.

Distances between squares are a measure for similarity between samples. A positive association of a gene with a given sample (i.e. an upregulation of that gene in that particular sample) results in the positioning of the gene and sample on a common line through the centroid (depicted by a cross). Correspondence analysis clearly identifies time as the major discriminator between the samples. In addition the effect of treatment with CRH can be identified as most prominent in the early time points.

Figure 3: A heat map depicting genes that are changed upon CRH treatment. Values were calculated by dividing the intensity of each sample by the intensity of the DMSO sample at the corresponding time point. These calculated ratio are converted into a color ramp using on a \log_2 scale. In this way the different timing of induction of expression becomes apparent. Genes showing a 2-fold change after 30 minutes of treatment with CRH were called "early responders", "intermediate responders" show a change after 1 to 2 hours of treatment and "late responders" show a response after 2 hours or more.

Figure 4: Overview of a selection of genes induced by CRH grouped by pathway or function as discussed in the text. Values were calculated by dividing the intensity of each sample by the intensity of the DMSO sample at the corresponding time point. These calculated ratio are converted into a color ramp using on a \log_2 scale and depicted in a heat map.

Figure 5: Induction of *Rgs2* by CRH in AtT-20 cells. Induction is calculated in comparison to levels observed in AtT-20 cells before any treatment. On top array data obtained for *Rgs2* are shown. Below, levels of *Rgs2* mRNA are shown as measured by quantitative RT-PCR on the same samples as used for array experiments and as measured on a repeated experiment.

30 DETAILED DESCRIPTION

As used herein, the term "compound" or "agent" means a biological or chemical compound such as a simple or complex organic molecule, a peptide, a protein or an oligonucleotide. A "test compound" as used herein, refers to a "compound" or "agent" used in a method according to the invention to assess whether said compounds modulates CRH signalling activity.

5 "CRH signaling" as used herein refers to the cellular changes in gene transcription after activation of the corticotropin releasing hormone receptor by CRH in said cell. It induces a CRH specific gene expression profile. Changes at the transcriptional level can be assessed either at the protein level or at the gene, RNA level.

10 "CRH response activity" as used herein refers in general to the change of a detectable cellular parameter as a result of the exposure of said cell to CRH. Detectable cellular parameters include amongst others, changes in membrane potential, changes in enzyme activity of an enzyme that modulates CRH signalling a said cell, changes in expression levels of a protein according to the invention or changes in the amount of second messengers such as cGMP, cAMP, Ca^{2+} or IP_3 .

15 The term "analog" or "functional analog" refers to a modified form of mammalian purine permeases in which at least one amino acid substitution has been made such that said analog retains substantially the same biological activity as the unmodified mammalian purine permease *in vivo* and/or *in vitro*.

20 The term "functional analog" is intended to include the "fragments," "variants," "degenerate variants," "analogs" and "homologues" or to "chemical derivatives" of the polypeptides according to the invention. Useful chemical derivatives of polypeptide are well known in the art and include, for example covalent modification of reactive organic site contained within the polypeptide with a secondary chemical moiety. Well known cross-linking reagents are useful to react to amino, carboxyl, or aldehyde residues to introduce, for example an affinity tag such as biotin, a fluorescent dye, or to conjugate the polypeptide to a solid phase surface (for example to create an affinity resin)

30 Variant(s) of polynucleotides or polypeptides, as the term is used herein, are polynucleotides or polypeptides that differ from a reference polynucleotide or polypeptide, respectively. A variant of the polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. (1) A polynucleotide that differs in nucleotide sequence from another, reference polynucleotide. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical. As noted below, changes in the nucleotide sequence of the variant may be silent. That is, they may not alter the amino acids encoded by the polynucleotide. Where alterations are limited to silent changes of this type a variant will encode a

polypeptide with the same amino acid sequence as the reference. Also as noted below, changes in the nucleotide sequence of the variant may alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Such nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed above. (2) A polypeptide that differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions, fusions and truncations, which may be present in any combination.

The terms "complementary" or "complementarity" as used herein refer to the capacity of purine and pyrimidine nucleotides to associate through hydrogen bonding to form double-stranded nucleic acid molecules. The following base pairs are related by complementarity: guanine and cytosine; adenine and thymine; and adenine and uracil. As used herein "complementary" means that the aforementioned relationship applies to substantially all base pairs comprising two single-stranded nucleic acid molecules over the entire length of said molecules. "Partially complementary" refers to the aforementioned relationship in which one of the two single-stranded nucleic acid molecules is shorter in length than the other such that a portion of one of the molecules remains single-stranded.

The term "conservative substitution" or "conservative amino acid substitution" refers to a replacement of one or more amino acid residue(s) in a parent protein without affecting the biological activity of the parent molecule based on the art recognized substitutability of certain amino acids (See e.g. M. Dayhoff, In Atlas of Protein Sequence and Structure, Vol. 5, Supp. 3, pgs 345-352, 1978).

"Fragment thereof" refers to a fragment, piece, or sub-region of a nucleic acid or protein molecule whose sequence is disclosed herein, such that said fragment comprises 5 or more amino acids, or 10 or more nucleotides that are contiguous in the parent protein or nucleic acid molecule.

"Functional fragment" as used herein, refers to an isolated sub-region, or fragment of a protein disclosed herein, or sequence of amino acids that, for example, comprises a functionally distinct region such as an active site for a receptor. Functional fragments may be produced by cloning technology, or as the natural products of alternative splicing mechanisms.

The term "homolog" or "homologous" describes the relationship between different nucleic acid molecules or amino acid sequences in which said sequences or molecules are related by partial identity or similarity at one or more blocks or regions within said molecules or sequences. "Isolated nucleic acid compound" refers to any RNA or DNA sequence, however construed or synthesized, which is locationally distinct from its natural location.

A "nucleic acid probe" or "probe" as used herein is a labeled nucleic acid compound which hybridizes with another nucleic acid compound. "Nucleic acid probe" means a single stranded nucleic acid sequence that will hybridize with a single stranded target nucleic acid sequence. A nucleic acid probe may be an oligonucleotide or a nucleotide polymer. A "probe" will usually contain a detectable moiety which may be attached to the end(s) of the probe or be internal to the sequence of the probe.

The term "primer" is a nucleic acid fragment which functions as an initiating substrate for enzymatic or synthetic elongation of, for example, a nucleic acid molecule.

The term "hybridization" as used herein refers to a process in which a single-stranded nucleic acid molecule joins with a complementary strand through nucleotide base pairing.

The term "stringency" refers to hybridization conditions. High stringency conditions disfavor non-homologous base pairing. Low stringency conditions have the opposite effect. Stringency may be altered, for example, by temperature and salt concentration. "Stringent conditions" refers to an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 x SSC at about 65°C. Further suitable hybridization conditions are described in the examples.

"Lower stringency conditions" include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1 X SSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC). Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's

reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

5 The term "fusion protein" as used herein refers to protein constructs that are the result of combining multiple protein domains or linker regions for the purpose of gaining the combined functions of the domains or linker regions. This may be accomplished by molecular cloning of the nucleotide sequences encoding such domains to produce a new polynucleotide sequence that encodes the desired fusion
10 protein. Alternatively, creation of a fusion protein may be accomplished by chemically joining two proteins.

 The term "linker region" or "linker domain" or similar such descriptive terms as used herein refers to polynucleotide or polypeptide sequence that are used in the construction of a cloning vector or fusion protein. Functions of a linker
15 region can include introduction of cloning sites into the nucleotide sequence, introduction of a flexible component or space-creating region between two protein domains, or creation of an affinity tag for specific molecule interaction. A linker region may be introduced into a fusion protein resulting from choices made during polypeptide or nucleotide sequence construction.

20

Screening methods

The present invention relates to screening methods to identify compounds that modulate corticotropin-releasing hormone (CRH) induced depression and stress. It is
25 based on the identification of a number of genes as downstream modulators of the CRH activated CRH receptors. In particular this invention provides a method for identifying a compound capable to alter the CRH signalling response in a cell, said method comprising;

- a) contacting said cell with CRH in the presence and absence of said compound;
 - 30 b) determine the change at transcriptional level of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said cell; and
 - c) compare the transcriptional level of said protein in the presence and absence of said compound;
- whereby the protein that modulates corticotropin releasing hormone (CRH)
35 signaling is being selected from the group consisting of SEQ ID NO.2, SEQ ID

4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID
NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ
ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32,
SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID
5 NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48.

To determine the change of transcription at the protein level one could
determine the amount of said protein using art known techniques. For example
using separation techniques such as isoelectric focusing or SDS-page in
10 combination with protein staining techniques such as coomassie or silver staining.
Alternatively, for proteins that are enzymes, the amount in a given solution or tissue
extract can be measured or assayed in terms of the catalytic effect the enzyme
produces, that is the conversion of its substrate into reaction product. For example,
for kinases one may assess the kinase activity using a substrate comprising the
15 kinase specific phosphorylation site and by measuring the phosphorylation of the
substrate by incorporation of radioactive phosphate into the substrate. This assay
may be performed both in the presence and absence of the compound to be tested.
For proteins that are not enzymes, other quantification methods are required. For
example transport proteins can be assayed by their binding to the molecule they
20 transport and hormones and toxins by the biological effect they produce.

To assess changes in transcription at the gene level, RNA or cDNA may be used
directly for detection or may be amplified enzymatically by using PCR or other
amplification techniques prior to analysis. Preferably said analysis method comprises
the use of a labelled oligonucleotide probe targeted to a suitable region of the gene.

25 Accordingly, in a preferred embodiment the level of gene transcription is
assessed using a probe which binds to a polynucleotide encoding an amino acid
sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID
NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16,
SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ
30 ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID
NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID
NO.48.

In another embodiment, an array of oligonucleotides probes comprising a
nucleotide sequence encoding a protein that modulates CRH signalling or fragments
35 thereof can be constructed to conduct efficient screening of gene expression. Array
technology methods are well known and have general applicability and can be used to
address a variety of questions in molecular genetics including gene expression, genetic

linkage, and genetic variability (see for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996)).

In an alternative embodiment, the method for identifying a compound capable to alter the CRH signalling response in a cell, comprises;

5 a) contacting said cell with CRH in the presence and absence of said compound;
 and

 b) determine the expression level of a polynucleotide comprising a nucleic acid
 sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO 3, SEQ ID
 NO.5, SEQ ID NO.7, SEQ ID NO.9, SEQ ID NO.11, SEQ ID NO.13, SEQ ID NO.15,
10 SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.25, SEQ
 ID NO.27, SEQ ID NO. 29, SEQ ID NO.31, SEQ ID NO.33, SEQ ID NO.35, SEQ ID
 NO.39, SEQ ID NO.41, SEQ ID NO.43, SEQ ID NO.45, SEQ ID NO.47 or SEQ ID
 NO:49.

To assess changes in expression levels, RNA or cDNA may be used directly for
15 detection or may be amplified enzymatically by using PCR or other amplification
 techniques prior to analysis. Preferably said analysis method comprises the use of a
 labelled oligonucleotide probe targeted to a suitable region of the polynucleotide.
 Accordingly, in a preferred embodiment the level of gene transcription is assessed using
 a probe which binds to a polynucleotide comprising a nucleic acid sequence selected
20 from the group consisting of SEQ ID NO: 1, SEQ ID NO 3, SEQ ID NO.5, SEQ ID
 NO.7, SEQ ID NO.9, SEQ ID NO.11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17,
 SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.25, SEQ ID NO.27, SEQ
 ID NO. 29, SEQ ID NO.31, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.39, SEQ ID
 NO.41, SEQ ID NO.43, SEQ ID NO.45, SEQ ID NO.47 or SEQ ID NO:49.

25 In another embodiment, an array of oligonucleotides probes comprising a
 nucleotide sequence encoding a protein that modulates CRH signalling or fragments
 thereof can be constructed to conduct efficient screening of gene expression.

 In another embodiment, an array of oligonucleotides probes comprising a
30 nucleotide sequence encoding a protein that modulates CRH signalling or fragments
 thereof can be constructed to conduct efficient screening of gene expression. Array
 technology methods are well known and have general applicability and can be used to
 address a variety of questions in molecular genetics including gene expression, genetic
 linkage, and genetic variability (see for example: M.Chee et al., Science, Vol 274, pp
35 610-613 (1996)).

 In case the genomic DNA is not used directly, mRNA may be isolated, and a
 first strand cDNA synthesis carried out. A second round of DNA synthesis can be

carried out for the production of the second strand. Subsequently by the specific PCR amplification an isolated cDNA can be obtained. If desired the double-stranded cDNA can be cloned into any suitable vector, for example, a plasmid, thereby forming a cDNA library. In analogy to the above, it is possible to screen
5 cDNA libraries constructed in a bacteriophage or plasmid shuttle vector with a labeled oligonucleotide probe targeted to any suitable region of the gene encoding a protein that modulates CRH signalling. See e.g. PCR Protocols: A Guide to Method and Application, Ed. M. Innis *et al.*, Academic Press (1990).

10 Methods for constructing cDNA libraries in a suitable vector such as a plasmid or phage for propagation in prokaryotic or eukaryotic cells are well known to those skilled in the art. [See e.g. Maniatis *et al. Supra*]. Suitable cloning vectors are well known and are widely available.

15 In a further embodiment changes in gene transcription are determined at mRNA level. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example; nucleic acid amplification, for instance via PCR, RT-PCR; RNase protection; Northern blotting and other hybridization
20 methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Assay techniques that can be used to determine the presence of protein derivatives or variants comprise amongst others mass spectrometry.

25 It is thus an object of the present invention to provide a method for identifying a compound capable to alter the CRH signalling response in a cell, said method comprising;

- a) contacting said cell with CRH in the presence and absence of said compound;
 - b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said cell; and
30
 - c) compare the amount of said protein in the presence and absence of said compound;
- whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.2, SEQ ID
35 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ

ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48.

5 Preferably, the method to assay the amount of protein that modulates CRH signaling is using an antibody which binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID NO.4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48.

15 Thus in another embodiment, this invention provides a monospecific antibody immunologically reactive with a protein that modulates CRH signalling said protein being selected from the group consisting of SEQ ID NO.2, SEQ ID NO.4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50. Antibodies generated against polypeptides of
20 the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells expressing these to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975)256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983)4:72) and the EBV-hybridoma technique (Cole *et al.*, *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, pp.77-96, Alan R. Liss, Inc., 1985).

30 Techniques for the production of single chain antibodies, such as those described in U.S. Patent No.4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity
35 chromatography.

Antibodies against polypeptides of the present invention may also be employed to treat the CRH metabolism related disorders such as CRH induced stress or

To determine the amount of protein that modulates CRH signalling, the antibodies according to the invention are used in conventional immunological techniques. Suitable immunological techniques are well known to those skilled in the art and include for example, ELISA, Western Blot analysis, competitive or sandwich immunoassays and the like, as is otherwise well known they all depend on the formation of an antigen-antibody immune complex wherein for the purpose of the assay, the antibody can be detectable labeled with, e.g. radio, enzyme or fluorescent labels or it can be immobilized on insoluble carriers.

For example in an ELISA screening format the antibody is added to a solid phase (for example the bottom of a microplate) which is coated with either the protein or a peptide fragment thereof coupled to a carrier (such as BSA), and then, adding an anti-immunoglobulin antibody (for example when the immunization is performed in mice, an anti-mouse immunoglobulin antibody is used, e.g. sheep-anti-mouse immunoglobulin (Ig)) conjugated with a detectable label such as an enzyme, preferably horseradish peroxidase, or a radioactive isotope such as ^{125}I .

It is thus an object of the invention to provide immunassays for the determination or detection of proteins that modulate CRH signalling in a sample, the method comprising contacting the sample with an antibody to the proteins according to the invention and determining whether an immune complex is formed between the antibody and said protein. These methods can either be performed on tissue samples or body fluid samples and generally comprise obtaining a sample from the body of a subject; contacting said sample with an imaging effective amount of a detectably labeled antibody according to the invention; and detecting the label to establish the presence of proteins that modulate CRH signalling in the sample.

The measuring methods using the antibodies of the present invention are not particularly limited. Any measuring method may be used as long as the amount of antibodies, antigens or the antigens-antibody complexes corresponding to the amount of the antigens to be measured is detected by chemical or physical means, and calculated from standard curves prepared by the use of standard solutions containing the antigens in known amounts. For example, nephelometry, competitive methods, immunometric methods and sandwich methods are suitably used. With respect to sensitivity and specificity, it is particularly preferred to use sandwich methods described below.

In measuring methods using labelling substances, radioisotopes, enzymes, fluorescent substances, luminous substances, etc. are used as labelling agents. Examples of the radioisotopes include ^{125}I , ^{131}I , ^3H and ^{14}C . Enzymes are usually made detectable by conjugation of an appropriate substrate that, in turn catalyzes a

detectable reaction. Examples thereof include, for example, beta-galactosidase, beta-glucosidase, alkaline phosphatase, peroxidase and malate dehydrogenase, preferably horseradish peroxidase. The luminous substances include, for example, luminol, luminol derivatives, luciferin, aequorin and luciferase. Further, the avidin-biotin systems can also be used for labelling the antibodies and immunogens of the present invention.

Accordingly, in a further aspect, the present invention provides for a method of identifying and obtaining compounds that alter the CRH signalling response activity in a cell, comprising:

a) contacting a cell which expresses at least one protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID NO.4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48, with said test compound; and

b) compare the CRH response activity of said cell in the presence and absence of said compound.

Changes in membrane potential can be measured using conventional electrophysiological techniques and when they become available, using novel high throughput methods currently under development. Since the change in membrane potential are normally the result of ion fluxes, as an alternative approach, changes in membrane potential can be measured indirectly through the change in intracellular ion concentrations using ion-sensitive fluorescent dyes, including fluo-3, fluo-4, fluo-5N, fura red, Sodium Green, SBFI and other similar probes from suppliers including Molecular Probes. Other fluorescent dyes, from suppliers including Molecular Probes, such as DIBAC₄(3) or Di-4-Anepps can detect membrane potential changes. For example calcium and sodium ion fluxes can thus be characterised in real time, using fluorometric and fluorescence imaging techniques, including fluorescence microscopy with or without laser confocal methods combined with image analysis algorithms.

In a preferred embodiment this assay is based around an instrument called a FLuorescence Imaging Plate Reader (FLIPR), Molecular Devices Corporation). -In its most common configuration, it excites and measures fluorescence emitted by

fluorescein-based dyes. It uses an argon-ion laser to produce high power excitation at 488 nm of a fluorophore, a system of optics to rapidly scan the over the bottom of a 96-/384-well plate and a sensitive, cooled CCD camera to capture the emitted fluorescence. It also contains a 96-/384-well pipetting head allowing the instrument to deliver
5 solutions of test agents into the wells of a 96-/384-well plate. The FLIPR assay is designed to measure fluorescence signals from populations of cells before, during and after addition of compounds, in real time, from all 96-/384-wells simultaneously.

It is thus an object of the present invention to provide a FLIPR assay used to screen for and characterise compounds functionally active in modulating CRH response
10 in cells, said cells expressing a protein selected from the group consisting of SEQ ID 26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32 and SEQ ID NO.34.

In an alternative embodiment, the activity of the cell may be assessed using electrophysiological methods. Therefore, proteins modulating CRH signalling in a cell can be characterised using whole cell and single channel electrophysiology.

15 It is thus a further object of this invention to provide a screening method to identify compounds which modulate CRH signalling response activity in a cell, said method comprising;

(a) contacting a host cell expressing a protein selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID
20 NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48, with a compound to be tested;

25 (b) measuring the effect of the test compound on the membrane potential of said cell using electrophysiological techniques; and

(c) compare the CRH response activity of said cell in the presence and absence of said compound.

30 In a preferred embodiment the host cells are *Xenopus* oocytes and the electrophysiological measurement consists of measuring the membrane current using the voltage clamp technique at distinct membrane potentials.

Changes in enzyme activity of an enzyme that modulates CRH signalling a said cell, can generally be measured or assayed in terms of the catalytic effect the enzyme produces, that is the conversion of its substrate into reaction product. For example, for
35 kinases one may assess the kinase activity using a substrate comprising the kinase specific phosphorylation site and by measuring the phosphorylation of the substrate. Similarly for phosphatases one may assess the phosphatase activity using a

phosphorylated substrate and by measuring the dephosphorylation of the substrate. These assays may be performed both in the presence and absence of the compound to be tested.

It is thus an object of the present invention to provide a method for identifying a compound capable of altering the CRH signalling response activity in a cell, said method comprising;

a) contacting a mixture comprising a kinase selected from the group consisting of SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16 and SEQ ID NO.18, with a source of phosphate and a suitable kinase substrate;

b) incubating said mixture in the presence or absence of a said compound and ; measuring the level of phosphorylation of said substrate in the presence of said compound compared to the level of phosphorylation of said substrate in the absence of said test compound.

In the assay of the invention, the kinase may be provided as a protein or it may be provided in the assay mixture as an mRNA encoding said kinase. When the assay comprises cell-free components, the kinase is provided as the protein. When the assay is conducted in the milieu of a cell, the kinase may be provided as either the protein or as an mRNA encoding said kinase, wherein, in order that the kinase be available in the assay, the mRNA is translated and kinase protein is thereby produced. It will be apparent from the Examples provided herein that it is a simple matter to obtain mRNA specifying the kinase and inject the mRNA into a cell for production of the kinase protein. The kinase may also be provided by expression of a plasmid, which encodes the kinase protein. Standard molecular biology techniques may be used to construct operable plasmids encoding the kinase protein and to express the plasmid in cells (Sambrook, et al., 1989, In: Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York).

As discussed herein, the method of identifying a kinase modulator may be performed either in vitro wherein the assay mixture is cell-free, in vitro wherein live cells are included in the assay, or in vivo in an animal. Thus, in one aspect of the invention, the mixture is contained within a eukaryotic cell and the method of the invention may be performed wherein some of the components of the assay mixture may be provided exogenously to a cell by microinjection of the components therein, and some of the components may be endogenous in the cell.

The term "endogenous in the cell" as used herein, means that the component is naturally produced in the subject cell.

The term "exogenous to the cell" as used herein, means that the component is not found naturally in the subject cell, or is found therein at a low level, and is added thereto.

When the method of the invention is performed using a eukaryotic cell, one or more of the kinase protein, the kinase substrate and the test compound may be injected into the eukaryotic cell prior to the incubation. The cell so injected is then incubated under conditions that facilitate protein kinase activity and the level of protein kinase activity is subsequently measured following the incubation period using the assays described herein.

The eukaryotic cell that is useful in the methods of the invention may be any one of a *Xenopus laevis* oocyte, a *Xenopus laevis* embryo cell, a mammalian cell (such as a I OTI/2 cell), a *Drosophila melanogaster* S2 cell, a *Dictyostelium discoideum* cell and a yeast cell. Still more preferably, the eukaryotic cell is the murine pituitary corticotroph-derived adenoma cell line cell AtT-20.

The source of phosphate for use in the methods of the invention may be any common source of phosphate, including, but not limited to, a nucleotide triphosphates such as, but not limited to, ATP or GTP. In a preferred embodiment, the phosphate source has bound thereon a detectable label which label is transferred with the phosphate group to the kinase substrate during the reaction. In this manner, phosphorylated kinase substrate may be distinguished from non-phosphorylated kinase substrate in that the phosphorylated substrate will contain the detectable label whereas the non-phosphorylated substrate will not contain the label. In another embodiment, the phosphate source does not have bound thereon a detectable label; instead, phosphorylated kinase substrate may be distinguished from non-phosphorylated kinase substrate, for instance by recognition of one form of the substrate, but not the other, by an antibody.

The detectable label which is useful in the methods of the invention may include any known or heretofore unknown detectable label which is transferred to the kinase substrate upon transfer of a phosphate group thereto as a result of protein kinase activity.

Labels which are useful include, but are not limited to, radioactive labels, such as $\gamma^{32}\text{P}$, ^{31}S , and non-radioactive labels, such as biotin and the like.

In another embodiment, the present invention to provide a method for identifying a compound capable to alter the CRH signalling response activity in a cell, said method comprising;

- a) contacting a mixture comprising a phosphatase selected from the group consisting of SEQ ID NO.36 and SEQ ID NO.38, and a suitable phosphorylated substrate;
- b) incubating said mixture in the presence or absence of a said compound and ;

measuring the level of phosphorylation of said substrate in the presence of said compound compared to the level of phosphorylation of said substrate in the absence of said test compound.

As for the kinase assay, the phosphatase in the assay of the invention, may be provided
5 as a protein or it may be provided in the assay mixture as an mRNA encoding said phosphatase. The phosphorylated substrate is typically labeled with a detectable phosphate residue. Labels which are useful include, but are not limited to, radioactive labels, such as $\gamma^{32}\text{P}$, ^{31}S , and non-radioactive labels, such as biotin and the like. For
10 use in a phosphatase activity assay, the substrate preferably consists of a peptide substrate, phosphorylated at a tyrosine or serine residue, typically labeled with $\gamma^{32}\text{P}$. In general, phosphorylation may be accomplished in a variety of ways. Typically, a protein tyrosine kinase is used. For example, a soluble EGF-receptor kinase in combination with $\gamma^{32}\text{P}$ -labeled ATP may be used to phosphorylate a tyrosine residue on a
15 peptide of the present invention. Such a phosphorylation reaction is typically allowed to proceed for about 2 hours at 30.degree. C. or overnight at room temperature. Phosphorylated peptide, hereinafter referred to as "phosphopeptide", is then purified from a phosphorylation reaction mixture. For example, peptide may be separated from a reaction mixture by addition of trichloroacetic acid and centrifugation, whereby the peptide remains in the supernate. The peptide is generally further purified by column
20 chromatography, e.g., on C18. Purified phosphorylated peptide may be lyophilized and stored at - 20.degree. C. prior to use.

Following incubation, phosphopeptide which is not dephosphorylated ("non-dephosphorylated phosphopeptide") is separated from radioactivity released by dephosphorylation of phosphopeptide (i.e., from free radioactive phosphorus released
25 by dephosphorylation). As used herein, the term "radioactive phosphorous" includes all forms in which a radioactive phosphorus atom may be present on a tyrosine residue and removed by dephosphorylation, e.g., as a phosphate group. Typically, separation of non-dephosphorylated phosphopeptide from free radioactive phosphorus released by dephosphorylation of phosphopeptide is effected by centrifugation, following
30 termination of the dephosphorylation reaction by the addition of substances including nonradioactive phosphates and charcoal. Radioactivity in the supernate is determined by means well known to those of ordinary skill in the art. Based upon the amount of radioactivity added to the assay mixture initially via the phosphopeptide and the amount of radioactivity detected at the end of the assay as radioactivity released by
35 dephosphorylation, the phosphatase enzymatic activity of the sample assayed may be calculated.



- 5

10

15

host cell for replication and/or appropriate expression of the exogenous DNA by the host cell. Accordingly, in a specific embodiment said vector is an expression vector such as pGL3luc, pBLCAT5 (LMBP 2451), pGMCSFlacZ (LMBP 2979), pEGFP or pSEAPbasic (DMB 3115), wherein LMBP and DMB numbers refer to the accession
5 numbers of these expression vectors at the Belgian Co-ordinated Collections of Micro-organisms.

In another aspect, the invention provides a method for identification of a compound modulating CRH signalling activity, said method comprising the steps: (i) contacting a
10 candidate agent with a gene promoter region as defined above; and (ii) determining whether said candidate agent modulates expression of the detectable product, such modulation being indicative for an agent capable of modulating CRH signalling activity. The detectable product refers either to the gene encoded protein or to the product of a reporter gene such as luciferase, β -galactosidase or green fluorescent
15 protein. Methods to quantify the detectable products are generally known in the art and include amongst others the use of a colorimetric substrate if the expression product is an enzyme, the use of specific antibodies in an RIA or ELISA assay or the measurement of the level of mRNA transcribed from genes operably linked to the promoter, wherein said mRNA can be measured either directly or indirectly using standard procedures.
20 Preferably, the gene promoter comprising a transcription factor binding site for a transcription factor selected from the group consisting of SEQ ID NO.2, SEQ ID NO.4, SEQ ID NO.6 and SEQ ID NO.8.

25 *Identification of the nucleic acid sequences encoding proteins capable of modulating CRH signaling.*

In another aspect the present invention relates to isolated and purified nucleic acid molecules which encodes proteins capable of modulating CRH signalling, wherein said
30 nucleic acid molecule is either RNA, DNA, cDNA or genomic DNA.

In particular, the present invention encompasses an isolated and purified nucleic acid molecule comprising a member selected from a group consisting of:

(a) a nucleic acid molecule encoding a protein that modulates CRH signalling
35 having at least a 70% identity to a polypeptide comprising an amino acid

sequence selected from the group consisting of SEQ ID No.46 and SEQ ID No.48 ;

(b) a nucleic acid molecule which is complementary to the polynucleotide of (a);

5 (c) a nucleic acid molecule comprising at least 15 sequential bases of the polynucleotide of (a) or (b);

(d) a nucleic acid molecule that hybridizes under stringent conditions to the polynucleotide molecule of (a) or (b); and

10 (e) a nucleic acid molecule encoding a protein that modulates CRH signalling comprising a nucleotide sequence of which is degenerated as a result of the genetic code to a nucleotide sequence of a polynucleotide of any of (a) to (d).

Those skilled in the art will recognize that owing to the degeneracy of the genetic code, numerous "silent" substitutions of nucleotide base pairs could be introduced into the sequences identified as SEQ ID NO:45, SEQ ID NO 47 or SEQ ID NO:49
15 without altering the identity of the encoded amino acid(s) or protein products. All such substitutions are intended to be within the scope of the invention.

In a further aspect, the present invention relates to Human purine permease polynucleotides. Such polynucleotides include isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide which has at least 70% identity,
20 preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to the amino acid sequence selected from the group consisting of SEQ ID NO:46 and SEQ ID NO:48, over the entire length said amino acid sequence. In this regard, polypeptides which have at least 97% identity are highly preferred, while those with at least 98-99% are more highly preferred, and
25 those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide consisting essentially of a polynucleotide sequence selected from SEQ ID No.45, SEQ ID No.47 or SEQ ID No.49.

Accordingly, in a further aspect, the present invention provides for an
30 isolated polynucleotide, comprising:

a) a nucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, more preferably at least 90% identity, yet more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity to an
35 amino acid sequence selected from the group consisting of SEQ ID No. 46 and SEQ ID No.48;

- b) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, more preferably at least 90% identity, Yet more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to a polynucleotide selected from the group consisting of SEQ ID No.45, SEQ ID No.47 and SEQ ID No.49, over the entire length of said polynucleotide;
- c) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, more preferably at least 90% identity, Yet more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity a polynucleotide selected from the group consisting of SEQ ID No.45, SEQ ID No.47 and SEQ ID No.49, over the entire coding region of said polynucleotide; and
- d) a nucleotide sequence consisting of a polynucleotide selected from the group consisting of SEQ ID No.45, SEQ ID No.47 and SEQ ID No.49.
- 15 The polynucleotides as outlined above are in particular provided for use in a screening method or diagnostic method according to the invention. In particular to identify capable of modulating CRH signaling in a cell or to diagnose altered CRH metabolism in an individual.

Identity or similarity, as known in the art, are relationships between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. Both identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity and similarity between two polynucleotide or two polypeptide sequences, both terms are well known to skilled artisans (Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., (1988) SIAM J. Applied Math., 48, 1073. Methods commonly employed to determine identity or similarity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., (1988)

The nucleic acid sequence encoding a protein capable of modulating CRH activity, or fragment thereof, can be isolated from a tissue in which said gene is expressed, such as but not limited to, brain, heart, kidney, pancreas, liver and skin. Said sequence can also be isolated from mammals other than human and mouse. Other cells and cell lines may also be suitable for use to isolate mammalian purine permease cDNA. Selection of suitable cells may be done by screening for CRH modulating activity in cell extracts or in whole cell assays, as described herein. Cells that possess CRH modulating activity in any one of these assays may be suitable for the isolation of purine permease DNA or mRNA.

35 Methods for constructing cDNA libraries in a suitable vector such as a plasmid or phage for propagation in prokaryotic or eukaryotic cells are well known to those skilled in the art. [See e.g. Maniatis *et al. Supra*]. Suitable cloning vectors are well known and are widely available.

It is readily apparent to those skilled in the art that other types of libraries, as well as libraries constructed from other cells or cell types, may be useful for isolating the nucleic acid sequences according to the invention. Other types of libraries include, but are not limited to, cDNA libraries derived from other cells, from organisms other than human and mouse, and genomic DNA libraries that include YAC (yeast artificial chromosome) and cosmid libraries. Construction of genomic DNA libraries can be performed by standard techniques well known in the art. Well known genomic DNA library construction techniques can be found in T. Maniatis *et al.* Molecular Cloning: A Laboratory Manual, 2d Ed. Chap. 14 (1989).

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide is short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low 'processivity' (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during the 1st strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (Frohman *et al.*, 1988, PNAS USA 85, 8998-9002), or recent modifications of this technique, exemplified by the MarathonTM technology (Clontech Laboratories Inc.).

In order to clone the polynucleotide encoding a protein according to the invention by the above methods, the amino acid sequence of the polypeptide encoded by said nucleic acid sequence may be necessary. To accomplish this, the proteins according to the invention, may be purified and partial amino acid sequence determined by automated sequencers. It is not necessary to determine the entire amino acid sequence, but the linear sequence of two regions of 6 to 8 amino acids from the protein is determined for the production of primers for PCR amplification of a partial DNA fragment.

Once suitable amino acid sequences have been identified, the DNA sequences capable of encoding them are synthesized. Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore, the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the polynucleotide sequences according to the invention and will be capable of hybridizing to DNA encoding the desired protein, even in the presence of DNA oligonucleotides with mismatches. DNA isolated by these methods can be used to

screen DNA libraries from a variety of cell types, from invertebrate and vertebrate sources, and to isolate homologous genes.

polypeptides

5 In a further embodiment this invention relates to a polypeptide in a substantially pure form which modulate CRH signalling wherein said polypeptide is encoded by an isolated and purified nucleic acid molecule according to the invention. In a preferred embodiment the polypeptide has the amino acid sequence selected from the group consisting of SEQ ID NO 46, SEQ ID NO 48 and
10 functional analogs thereof

The protein according to the invention includes all possible amino acid variants encoded by the nucleic acid according to the invention including a polypeptide encoded by said molecule and having conservative amino acid changes.

Those skilled in the art will recognize that the protein which modulate CRH
15 signalling could be obtained by a plurality of recombinant DNA techniques including, for example, hybridization, polymerase chain reaction (PCR) amplification, or *de novo* DNA synthesis (See e.g., T. Maniatis *et al.* Molecular Cloning: A Laboratory Manual, 2d Ed. Chap. 14 (1989)).

Purified biologically active protein that modulates CRH signalings may have
20 several different physical forms. The polypeptides according to the invention may exist as full-length nascent or unprocessed polypeptides, or as partially processed polypeptides or combinations of processed polypeptides. The full-length nascent polypeptide may be post-translationally modified, amongst other, by specific proteolytic cleavage events that result in the formation of fragments of the full-length
25 nascent polypeptide. A fragment, or physical association of fragments may have the full biological activity associated with proteins according to the invention; however, the degree of CRH modulating activity may vary between individual fragments.

Also preferred in this aspect of the invention are fragments characterized by structural or functional attributes of the polypeptide. Preferred embodiments of the
30 invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, high antigenic index regions of the
35 polypeptide of the invention, and combinations of such fragments. Preferred regions are those that mediate activities of the polypeptides of the invention. Most highly preferred in this regard are fragments that have a chemical, biological or other

activity of the response regulator polypeptide of the invention, including those with a similar activity or an improved activity, or with a decreased undesirable activity.

Recombinant expression of polynucleotides encoding protein which modulate CRH activity

In another embodiment polynucleotides according to the invention may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce a protein that modulates CRH signalling. Techniques for such manipulations are fully described in Maniatis, T, et al., supra, and are well known in the art.

Therefore, in a further aspect this invention provides an expression vector for expression of a protein that modulates CRH signalling in a recombinant host, wherein said vector contains a nucleic acid sequence encoding a protein that modulates CRH signalling and functional analogs thereof. In a more preferred aspect of this invention this expression vector contains a nucleic acid molecule encoding a protein that modulates CRH signalling, having a nucleotide sequence selected from a group consisting of: SEQ ID NO:45, SEQ ID NO 47, SEQ ID NO:49 and functional analogs thereof or contains genomic DNA encoding a protein that modulates CRH signalling.

Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic genes in a variety of hosts such as bacteria including E. coli, cyanobacteria, plant cells, insect cells, amphibian cells, fungal cells including yeast cells, and animal cells.

Specifically designed vectors allow the shuttling of DNA between hosts such as bacteria-yeast or bacteria-animal cells or bacteria-fungal cells or bacteria-invertebrate cells. An appropriately constructed expression vector may contain: an origin of replication for autonomous replication in host cells, selectable markers, a limited number of useful restriction enzyme sites, a potential for high copy number, and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA synthesis. A strong promoter is one that causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

The isolated and purified nucleic acid molecules, according to the invention, encoding a protein which modulates CRH signalling may be cloned into an expression vector for expression in a recombinant host cell. Recombinant host cells may be prokaryotic or eukaryotic, including but not limited to bacteria such as E. coli, fungal cells such as yeast, amphibian cells such as Xenopus oocytes, mammalian cells including but not limited to cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not limited to Drosophila- and silkworm-derived cell lines. Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26), MRC-5 (ATCC CCL 171), L-cells, neuroblastoma, glial cells and HEK-293 (ATCC CRL1573).

Therefore, in a further embodiment this invention relates to a recombinant host cell containing a recombinantly cloned nucleic acid molecule encoding a protein that modulates CRH signalling or functional analog thereof. In a further aspect the recombinant host cell according to the invention contains a nucleic acid molecule which is either genomic DNA or has a nucleotide sequence selected from a group consisting of: (SEQ ID NO:45); (SEQ ID NO 47); (SEQ ID NO:49); and functional analogs thereof.

The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, protoplast fusion, lipofection, and electroporation. The expression vector-containing cells are clonally propagated and analyzed to determine whether they produce a protein that modulates CRH signalling. Identification of permeases expressing host cell clones may be done by several means, including but not limited to immunological reactivity with antibodies directed against the polypeptides according to the invention, and the presence of host cell-associated mammalian purine permease activity.

Thus, the present invention also relates to a process for expression of protein that modulates CRH signalling in a recombinant host cell, comprising culturing the host cells according to the invention under conditions which allow expression of the protein that modulates CRH signalling from the expression vector as outlined herein. The proteins of this invention may be synthesized either by direct expression or as a fusion protein comprising the protein of interest as a translational fusion with another protein or peptide that may be removable by self, enzymatic or

chemical cleavage. Therefore, in a particular embodiment this invention provides the proteins according to the invention wherein said polypeptides are part of a fusion protein.

5 It is often observed in the production of certain peptides in recombinant systems that expression as a fusion protein prolongs the life span, increases the yield of the desired peptide, or provides a convenient means of purifying the protein. This is particularly relevant when expressing mammalian proteins in prokaryotic hosts. A variety of peptidases (e.g. enterokinase and thrombin), which
10 cleave a polypeptide at specific sites or digest, the peptides from the amino or carboxy termini (e.g. diaminopeptidase) of the peptide chain are known. Furthermore, particular chemicals (e.g. cyanogen bromide) will cleave a polypeptide chain at specific sites. The skilled artisan will appreciate the modifications necessary to the amino acid sequence (and synthetic or semi-synthetic coding sequence if recombinant means are employed) to incorporate site-specific
15 internal cleavage sites. See e.g., P.Carter, "Site Specific Proteolysis of Fusion Proteins", Chapter 13, in Protein Purification: From Molecular Mechanisms to Large Scale Processes, American Chemical Society, Washington, D.C. (1990).

Furthermore, one could use, e.g., a mammalian cell that already comprises in its genome a nucleic acid molecule encoding a protein that modulates CRH
20 signalling as described above, but does not express the same or not in an appropriate manner due to, e.g., a weak promoter, and introduce into the mammalian cell a regulatory sequence such as a strong promoter in close proximity to the endogenous nucleic acid molecule encoding said purine permease polypeptide so as to induce expression of the same.

25 As such a recombinant host cell containing a polynucleotide encoding a protein which modulates CRH signalling under the control of a heterologous transcription and/or regulatory sequence or protein, would be another embodiment of this invention.

In this context the term "regulatory sequence" denotes a nucleic acid
30 molecule that can be used to increase the expression of the purine permease polypeptide, due to its integration into the genome of a cell in close proximity to the CRH modulating protein-encoding gene. Such regulatory sequences comprise promoters, enhancers, inactivated silencer intron sequences, 3'UTR and/or 5'UTR coding regions, protein and/or RNA stabilizing elements, nucleic acid molecules
35 encoding a regulatory protein, e.g., a transcription factor, capable of inducing or triggering the expression of the CRH modulating protein-encoding gene or other gene expression control elements which are known to activate gene expression

and/or increase the amount of the gene product. The introduction of said regulatory sequence leads to increase and/or induction of expression of polypeptides, which modulate CRH signalling, resulting in the end in an increased amount of said polypeptides in the cell. Thus, the present invention is aiming at providing *de novo* and/or increased expression of polypeptides that modulate CRH signalling.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis, Basic Methods In Molecular Biology (1986). It is specifically contemplated that polypeptides, which modulate CRH signalling may in fact be expressed by a host cell lacking a recombinant vector.

In addition, expression of polynucleotides according to the invention may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA or mRNA isolated from cells capable of modulating CRH signaling can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based systems, including but not limited to microinjection into frog oocytes, with microinjection into frog oocytes being generally preferred.

Transgenic non-human animals

The present invention also relates to a method for the production of a transgenic non-human animal, preferably transgenic mouse, comprising introduction of a polynucleotide or vector of the invention into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with a screening method of the invention described herein and may be a non-transgenic healthy animal, or may have a phosphate uptake or reabsorption disorder, preferably a disorder caused by at least one mutation in the protein that modulates CRH signalling. Such transgenic animals are well suited for, e.g., pharmacological studies of drugs in connection with mutant forms of the above described polypeptides. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., Gene Targeting, A Practical Approach (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using, e.g., Southern blots with an appropriate probe; see *supra*.

15

20

25

35

5

15

- 20

30

It will thus be appreciated that the present invention also relates to a method for preventing, treating or ameliorating a medical condition related to a disorder of CRH metabolism including CRH receptor related disorders which comprises administering to a mammalian subject a therapeutically effective amount of the polypeptides, the polynucleotides or the vectors encoding a protein capable of modulating CRH signalling of the present invention.

Diagnostic Assays

This invention further relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO: 1, SEQ ID NO 3, SEQ ID NO.5, SEQ ID NO.7, SEQ ID NO.9, SEQ ID NO.11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.25, SEQ ID NO.27, SEQ ID NO. 29, SEQ ID NO.31, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.39, SEQ ID NO.41, SEQ ID NO.43, SEQ ID NO.45, SEQ ID NO.47 or SEQ ID NO:49 which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

It will thus be appreciated that this invention provides a method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to a disorder of CRH activity comprising:

(a) determining the presence or absence of a mutation in the polynucleotide according to the invention; and

(b) diagnosing a pathological condition or susceptibility to a pathological condition based on the presence or absence of said mutation.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled mammalian purine permease nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures.

DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in capillary electrophoresis columns or gels, with or without denaturing agents, or by direct DNA sequencing (e.g., Myers *et al.*, *Science* (1985)230:1242). Sequence changes at specific locations may also be revealed by
5 specific restriction endonucleases, nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method (see Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85: 4397-4401). In another embodiment, an array of oligonucleotides probes comprising a nucleotide sequence encoding a protein capable of modulating CRH
10 activity or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

The diagnostic assays offer a process for diagnosing or determining a
15 susceptibility to the diseases through detection of mutations in the CRH modulating protein-encoding gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of polypeptide or mRNA, as well as by determining from said samples the presence of protein derivatives compared to the
20 normal structure. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example; nucleic acid amplification, for instance via PCR, RT-PCR; RNase protection; Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as
25 a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Assay techniques that can be used to determine the presence of protein derivatives or variants comprise amongst others mass spectrometry.

30 Thus in another aspect, the present invention provides a method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to a disorder of prolonged CRH exposure comprising:

- (a) determining the presence or amount of expression of the polypeptide or a derivative thereof according to the invention in a biological sample; and
- 35 (b) diagnosing a pathological condition or susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide or of a derivative thereof.

In particular the present invention provides a method of diagnosing a CRH induced gene expression profile in an individual, said method comprising;

a) obtaining a biological sample of said individual; and

b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said biological sample;

whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48

Preferably the amount of said proteins is determined either at the protein level, preferably using antibodies that bind thereto, or at the gene transcription level, preferably using probes that bind to a polynucleotide encoding an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46 and SEQ ID NO.48

Alternatively the CRH induced gene expression profile is determined by assessing the level of gene transcription of a gene comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO 3, SEQ ID NO.5, SEQ ID NO.7, SEQ ID NO.9, SEQ ID NO.11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.25, SEQ ID NO.27, SEQ ID NO. 29, SEQ ID NO.31, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.39, SEQ ID NO.41, SEQ ID NO.43, SEQ ID NO.45, SEQ ID NO.47 or SEQ ID NO:49. Methods to determine the level of gene transcription have been described hereinbefore and comprise in a preferred embodiment the use of a probe which binds, preferably selectively binds to a polynucleotide selected from the group consisting of SEQ ID NO: 1, SEQ ID NO 3, SEQ ID NO.5, SEQ ID NO.7, SEQ ID NO.9, SEQ ID NO.11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.25, SEQ ID NO.27, SEQ ID

NO. 29, SEQ ID NO.31, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.39, SEQ ID NO.41, SEQ ID NO.43, SEQ ID NO.45, SEQ ID NO.47 or SEQ ID NO:49 or the complement thereof. . In another embodiment, an array of oligonucleotides probes comprising a nucleotide sequence according to the invention or fragments thereof
5 can be constructed to conduct efficient screening of the level of gene transcription in the sample of an individual.

In a further aspect, the present invention relates to a diagnostic kit which comprises:

- 10 (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 51, SEQ ID NO: 53, SEQ ID NO: 55 or a fragment thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO 56 or a fragment thereof; or
- 15 (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of NO:52, SEQ ID NO:54 or SEQ ID NO 56 and optionally suitable means for detection.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or
20 susceptibility to a disease, particularly CRH metabolism related disorders such as CRH induced stress or depression.

The nucleotide sequences of the present invention are also valuable for chromosome localisation. These sequences are specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The
25 mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene-associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in
30 Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). The gene of the present invention maps to human chromosome 15.

35 The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is

likely to be the causative agent of the disease.

The nucleotide sequences of the present invention are also valuable for tissue localisation. Such techniques allow the determination of expression patterns of the polypeptides according to the invention in tissues by detection of the mRNAs that
5 encode them. These techniques include *in situ* hybridization techniques and nucleotide amplification techniques, for example PCR. Such techniques are well known in the art. Results from these studies provide an indication of the normal functions of the polypeptides in the organism.

10 The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them, can also be used as immunogens to produce antibodies immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in
15 the prior art.

Thus in another embodiment, this invention provides a monospecific antibody immunologically reactive with a mammalian purine permease. In a preferred embodiment said antibody is immunologically reactive with a polypeptide having an amino acid sequence selected from a group consisting of: (SEQ ID NO:52); (SEQ ID
20 NO 54); (SEQ ID NO:56); and functional analogs thereof or said antibody blocks activity of a protein that modulates CRH signalling.

Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells expressing these to an animal, preferably a non-human animal, using routine
25 protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975)256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983)4:72) and the EBV-hybridoma technique (Cole *et al.*,
30 MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp.77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No.4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other
35 organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity

chromatography.

Antibodies against polypeptides of the present invention may also be employed to treat the CRH metabolism related disorders.

5 In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgD, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG I, where fusion takes place at the hinge region. In a particular
10 embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with for instance blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides
15 encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. W094/29458 and W094/22914.

Therapeutic Utility

A further aspect of the invention relates to an immunological/vaccine
20 formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a polypeptide of the present invention wherein the composition comprises a polypeptide or polynucleotide of the present invention. The vaccine formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered
25 parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include
30 suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific
35 activity of the vaccine and can be readily determined by routine experimentation.

In still another approach, expression of the gene encoding proteins which modulate CRH signalling can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally
5 generated or externally administered (see, for example, O'Connor, *J.Neurochem* (1991) 56:560 ;Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices ("triplexes") with the gene can be supplied (see, for example, Lee *et al.*,
10 *Nucleic Acids Res* (1979)6:3073; Cooney *et al.*, *Science* (1988)241:456; Dervan *et al.*, *Science* (1991)251:1360). These oligomers can be administered per se or the relevant oligomers can be expressed *in vivo*. Synthetic antisense or triplex oligonucleotides may comprise modified bases or modified backbones. Examples of the latter include methylphosphonate, phosphorothioate or peptide nucleic acid
15 backbones. Such backbones are incorporated in the antisense or triplex oligonucleotide in order to provide protection from degradation by nucleases and are well known in the art. Antisense and triplex molecules synthesised with these and/or other modified backbones also form part of the present invention.

In another process for inhibiting expression of a target gene in a cell, RNA with partial or fully double-stranded character is introduced into the cell or into the
20 extracellular environment. Inhibition is specific in that a nucleotide sequence from a portion of the target gene is chosen to produce inhibitory RNA. The RNA may comprise one or more strands of polymerized ribonucleotide; it may include modifications to either the phosphate-sugar backbone or the nucleoside. The double-stranded structure may be formed by a single self-complementary RNA strand or two
25 complementary strands. Inhibition is sequence-specific in that the nucleotide sequences corresponding to the duplex region of the RNA are targeted for genetic inhibition. RNA containing a nucleotide sequence identical to a portion of the target sequence is preferred. Examples of RNA inhibition technology can be found in International Patent Application WO 99/32619.

30 In addition, expression of the proteins which modulate CRH signalling may be prevented by using ribozymes specific to the mRNA sequence encoding said protein. Ribozymes are catalytically active RNAs that can be natural or synthetic (see for example Usman, N, *et al.*, *Curr. Opin. Struct. Biol* (1996)6(4), 527-33.) Synthetic
35 ribozymes can be designed to specifically cleave the aforementioned mRNAs at selected positions thereby preventing translation of said mRNAs into functional polypeptide. Ribozymes may be synthesised with a natural ribose phosphate backbone and natural bases, as normally found in RNA molecules. Alternatively the

ribozymes may be synthesised with non-natural backbones to provide protection from ribonuclease degradation, for example, 2'-O-methyl RNA, and may contain modified bases.

For treating abnormal conditions related to an under-expression of proteins which modulate CRH signalling, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates a polypeptide of the present invention, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of mammalian purine permease by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication-defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of a polypeptide of the present invention in combination with a suitable pharmaceutical carrier.

In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide, such as the soluble form of a polypeptide of the present invention, agonist/antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as

subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of patches, salves, pastes, gels, and the like.

The dosage range required depends on the choice of peptide or other compounds of the present invention, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 μ g/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

This invention will be better understood by reference to the Experimental Details that follow, but those skilled in the art will readily appreciate that these are only illustrative of the invention as described more fully in the claims that follow thereafter. Additionally, throughout this application, various publications are cited. The disclosure of these publications is hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains.

EXPERIMENTAL PROCEDURES

Cell culture and sample preparation - AtT-20 cells were purchased from ATCC and were maintained at 37°C in 5% CO₂ in humidified air in Dulbecco's modified Eagle's medium (Invitrogen Life Technologies) containing 10 % fetal bovine serum, 5% horse

serum and 4.5 g/L D-glucose. For experiments cells were seeded in 25 cm² flasks. The medium was replaced 48 h later and cells were treated with either 0.1% DMSO, 1μM CRH (Sigma) in DMSO, 1μM R121919 in DMSO or 1μM CRH +1μM R121919 in DMSO for 0, 0.5, 1, 2, 4, 8 and 24 h in fresh medium. The incubation was stopped by aspirating the incubation medium and adding 3 ml Trizol (Invitrogen Life Technologies) for lysis of the cells. Total RNA was extracted using Trizol according to the instructions of the manufacturer. 100μg total RNA was further purified using Rneasy kit (Qiagen) with DNaseI treatment on column.

Microarray hybridization – cRNA was prepared as follows. Reverse transcription was performed on 10μg of total RNA for 1h at 42°C using a T7-oligo(dT)₂₄-primer and SuperscriptII RT (Invitrogen Life Technologies). Second strand cDNA synthesis was done for 2 h at 16°C using *Escherichia coli* DNA Polymerase I, DNA ligase and RNaseH (Invitrogen Life Technologies). After phenol-chloroform extraction using phase-lock gel (Eppendorf) in vitro transcription was performed for 6 h at 37°C using the Bioarray high-yield RNA transcript labeling kit with Biotin labeled ribonucleotides (Enzo Diagnostics). cRNA samples were purified on Qiagen Rneasy columns followed by fragmentation for 35 min at 95°C. cRNA yields were between 50 and 100μg. Samples were processed on GeneChips (Affymetrix, Santa Clara, CA). In order to check the quality of each sample, 5μg of labeled cRNA was run on Test2-arrays. Actual experiments were performed on Murine Genome U74Av2 arrays, containing probe sets interrogating approximately 12,000 full-length mouse genes and EST clusters from the UniGene database (Build 74). Hybridization was performed using 15μg cRNA for 16 h at 45°C under continuous rotation. Arrays were stained in Affymetrix Fluidics stations using Streptavidin/Phycoerythrin (SAPE) followed by staining with anti-streptavidin antibody and a second SAPE staining. Subsequently arrays were scanned with a HP-Laserscanner and data were analyzed with the Microarray Suite Software (Affymetrix). No scaling or normalization was performed at this stage. Quality of the experiment was assessed based on the percentages present calls across all samples which was on average 47.06±2.45%. The cytoplasmic β-actin and GAPDH 5'/3' ratios were 1.10±0.08 and 0.93±0.05 respectively.

Data analysis and selection of genes

Raw intensities on each array were aligned using the following global mean algorithm:

$$\log I_{aligned} = \log I_{raw} + \left(\log \frac{\sum_{all\ samples} I_{row}}{\# genes \times \# samples} - \log \frac{\sum_{sample} I_{row}}{\# genes} \right)$$

Basically this alignment sets the average intensity of one array to the average measured across all arrays, compensating for array to array variations in hybridization, washing and staining, ultimately allowing a reasonable comparison between arrays. After alignment polished data were analyzed using weighted spectral mapping. Weighted spectral mapping is an unsupervised multivariate analysis method which includes double-centering of the data combined with a specialized visualization representing the two highest principle components. Even though double-centering removes the "size" component of the array data, this information is reintroduced in the visualization via the area of the symbols representing the size of the respective samples and genes. This method allows the reduction of a large microarray dataset and provides means to visually inspect and thereby identify clusters of genes and/or subjects in the data (7). A more detailed analysis was carried out using the OmniViz program. All records which were absent in all experiments were removed and all signals less than 20 were set to 20. Gene expression fold differences for CRH, R121919 and CRH+R121919 treatments were calculated at each timepoint. For those calculations signals at corresponding timepoints in DMSO treated samples were used to calculate ratios.

Quantitative RT-PCR - Microarray data were confirmed using real time PCR analysis. First strand cDNA synthesis was performed on 0.5µg total RNA using random hexamer primers and SuperscriptII RT (Invitrogen Life Technologies). Quantitative PCR was performed on a ABIPrism 7700 cycler (Applied Biosystems) using a Taqman PCR kit. Serial dilutions of cDNA were used to generate standard curves of threshold cycles versus the logarithms of concentration for β -actin, *c-fos*, *Crh-R1*, *Crh-R2* and *Rgs2* (see table for sequences). A linear regression line calculated from the standard curves allowed the determination of transcript levels in RNA samples from the different time points.

β -actin forward	5'-CATCTTGGCCTCACTGTCCAC-3'
β -actin probe	5'-TGCTTGCTGATCCACATCTGCTGGA-3'
β -actin reverse	5'-GGGCCGGACTCATCGTACT-3'
c-fos forward	5'-GGGAGGACCTTACCTGTTTCGT-3'
c-fos probe	5'-CACCAGGCTGTGGGCCTCAAGG-3'
c-fos reverse	5'-CCAGATGTGGATGCTTGCAA-3'
Crh-R2 forward	5'-GGGAGAACAGAAGCGCCTG-3'
Crh-R2 probe	5'-AGAAGGGTGAGGATCCCCCAAATCAGAGT-3'
Crh-R2 reverse	5'-CCCTTGTTTCAATCACTCCCA-3'
Crh-R1 forward	5'-TTTCTGAACAGTGAGGTCCGC-3'

Crh-R1 probe	5'-CCGGAAGAGGTGGCGGCGA-3'
Crh-R1 reverse	5'-GGGCTCTGATGGAGTGCTTG-3'
Rgs2 forward	5'-TTGGAAGACCCGTTTGAGCTA-3'
Rgs2 probe	5'-TCTTGCAGAATTCCTCTGCTCCTGGG-3'
Rgs2 reverse	5'-TTTCTTGCCAGTTTGGGCT-3'

RESULTS

Transcriptional response to CRH was studied in the CRH-R1-expressing murine AtT-20 pituitary corticotroph-derived adenoma cell line. Whereas CRH-R1 was readily detectable both by real time quantitative RT-PCR (RTq) and western blot, CRH-R2 expression could not be discerned in AtT-20 cells. In order to identify CRH-R1 specific responses cells were exposed to 1 μ M CRH, 1 μ M CRH in the presence of 1 μ M of a CRH-R1 specific antagonist R121919 and to R121919 alone. Transcriptional responses were followed over time until 24 h after the first administration. In order to assess treatment efficacy, *c-fos* mRNA levels were determined by RTq on RNA from the different treatments and time points before array experiments were carried out. In agreement with previous reports, exposure to CRH elicited a transient surge in *c-fos* transcription, with levels already going down after 0.5 to 1 h (see figure 1)(8;9). This response was almost completely suppressed in the presence of R121919. Interestingly, 0.1% DMSO induced *c-fos* expression; however levels were between 5 to 10 times lower compared to CRH induced expression.

All time points were analyzed on microarrays containing approximately 12000 murine genes and ESTs. Overall analysis of the expression profiles using spectral mapping (a so called unsupervised method) indicated progressing time to account for most of the observed changes in gene expression (see figure 2). Synchronization of cell cycle in these cultures induced by addition of new medium and serum could possibly account for this phenomenon although accumulation of metabolites and progressing cell culture are additional contributing factors. The spectral map analysis also showed that CRH treated samples differed from other samples mainly in the early time points (0.5h until 2h), with overall differences in expression becoming very small after 8h. Because of this obvious influence of time, expression measurements were analyzed relative to those observed in the corresponding time point in DMSO treated control samples.

Regulated genes were defined as those showing a greater than 2-fold change in transcript levels at any one time point. Using the OmniViz Treescape view, 111 genes that met this criterion, showing a difference in expression after treatment with CRH compared to treatment with the antagonist, were selected. 26 out of the 111 genes were "early responders", showing a 2-fold change already after 30 minutes treatment with

CRH. 32 genes were "intermediate responders", responding after 1 to 2 hours of treatment and 53 genes were "late responders", showing a response after 2 hours or more after treatment (see figure 3). These responses were suppressed by the CRH-R1 antagonist R121919. Among the early responders were known players in the pathways downstream of the CRH-R1 such as the transcription factors *Nurr1*, *Nurr77*, *Jun-B*, validating the assay.

Interesting novel players identified include transcription factors (e.g. hairy/enhancer-of-split related 1 (*Hey1*), nuclear factor regulated by interleukin 3 (NFIL3), cAMP responsive element modulator (*CREM*) and prostate specific ets transcription factor (*Pse*)), receptor and channel regulators (e.g. Ras-related GTP-binding protein (*GEM*) and receptor (calcitonin) activity modifying protein 3 (*RAMP3*)), secreted peptides (e.g. adrenomedullin, calcitonin, cholecystokinin) and proteins involved in intracellular signaling (e.g. regulator of G-protein signaling 2 (*Rgs2*), cAMP specific phosphodiesterase 4B (*Pde4b*), inositol 1,4,5-triphosphate receptor 1 (*IP₃R1*) and the regulatory subunit phosphatidylinositol 3-kinase, p85). Other interesting regulated genes comprise Period homolog *Per1*, fibroblast growth factor receptor 2 (*Fgfr2*), serum/glucocorticoid regulated kinase and serum-inducible kinase (figure 4). Interestingly, all responders identified according to above mentioned criteria were up regulated after exposure to CRH. This induction was transient and nearly all of the induced genes return to baseline after 4 to 8 hr. Many of the induced transcripts encode proteins that would exert a negative feedback on the CRH-R1 signaling (e.g. *Pde4*, *Rgs2*, *CREM*, etc...).

Confirmation of microarray data was carried out using quantitative real time PCR analysis on the same samples used for hybridization experiments and on a repeated experiment. Levels of regulation identified by microarray corresponded to those observed by quantitative PCR as shown for *Rgs2* in figure 5.

DISCUSSION

We have identified transcriptional pathways downstream of the CRH-R1 in Art-20 cells using a CRH-R1 specific antagonist. Our findings are in agreement with activation of several second messenger such as cAMP and Ca^{2+} upon stimulation with CRH. Some of the transcriptional responses can be explained by the phosphorylation of CREB and the subsequent transcription of genes downstream of cAMP responsive elements. These elements have been found for example in the promoters of *Per1*, *Nurr1*, *CREM-ICER*, *c-Fos*. Furthermore the kinetic profile of the induction of these genes correspond with the observed maximal transcription rate by CREB after 0.5 hr of

cAMP formation. The induction of CREM-ICER constitutes a negative feedback mechanism in attenuating transcriptional response to cAMP. Of interest is the reported induction of CREM-ICER in response to acute stress in the intermediate lobe of the pituitary gland. Mice deficient for CREM-ICER show a chronic increase of beta-

5 endorphin levels suggesting that CREM-ICER induction may be involved in the modulation of gene expression in response to stress (10). Our results suggest that CREM-ICER is directly involved in the modulation of CRH signaling and as a result, ablation of CREM-ICER could lead to an altered response to stress signals. Another novel putative negative feedback regulator of CRH signaling is Rgs2. We identified

10 two single CRE motifs in the promoter of the human *RGS2* gene, providing a possible explanation for the early response behavior of this gene upon stimulation with CRH. In support of our findings is a recent report showing that both phosphoinositide signaling and cAMP induce a rapid and transient increase in Rgs2 mRNA in human astrocytoma and neuroblastoma cells. The Rgs2 protein is a selective inhibitor of G_q function.

15 Recently it has been shown that Rgs2 reduces odorant-elicited cAMP production, not by acting on G_q but by directly inhibiting the activity of adenylyl cyclase type III. Although Rgs2 was originally identified as an immediate early response gene in activated T lymphocytes, studies in Rgs2 deficient mice indicate that it also plays a role in the modulation of stress related behavior as these mice show increased anxiety and

20 aggression (11). Also the induction of cAMP specific phosphodiesterase 4B (*Pde4b*) can be categorized under negative feedback, directly attenuating the cAMP signal. Another important second messenger generated upon stimulation with CRH is Ca^{2+} . It has been shown that CRH triggers a steady-state depolarization stimulated extracellular Ca^{2+} entry via voltage-gated Ca^{2+} channels and raises intracellular Ca^{2+} concentration

25 through release from inositol 1,4,5-triphosphate (InsP3) sensitive Ca^{2+} pools(12). Both the InsP3 receptor and the p85 regulatory subunit of phosphatidylinositol 3-kinase are upregulated, possibly accounting for a compensating mechanism for prolonged Ca^{2+} signaling. Also the upregulation of the small G-protein kir/Gem points towards an attenuation of prolonged Ca^{2+} signaling. Recent studies have shown that Gem regulates

30 Ca^{2+} channel expression at the cell membrane through the β auxiliary subunits. Increased levels of Gem have been shown to inhibit Ca^{2+} -triggered exocytosis and it has been proposed that Gem could have a protective effect against Ca^{2+} overload. In addition to its role as second messenger, intracellular Ca^{2+} has also been shown to play a critical role in regulating gene expression. Of interest is the regulation of

35 NFIL3/E4BP4 by calcineurin/NFAT and CaM kinase signaling, accounting for an increase in NFIL3 mRNA levels upon CRH treatment. In B lymphocytes expression of NFIL3 is induced by interleukin 3 through both the Raf-mitogen-activated protein

kinase and phosphatidylinositol 3-kinase pathways. In this cell type NFIL3 inhibits apoptosis in synergy with Bcl-xL dependent pathways. Our data suggest a role for NFIL3 in prevention of apoptosis in AtT-20 cells.

5 CRH is the most efficacious ACTH secretagogue. Unfortunately the microarrays that were used did not interrogate for POMC levels. However several other prepropeptides
10 mRNAs were found upregulated after CRH administration such as cholecystokinin (CCK) and two calcitonin peptide family members, adrenomedullin (ADM) and calcitonin (CT). Also of interest in this respect is the upregulation of RAMP3. RAMPs control the transport and glycosylation of the calcitonin receptor-like receptor (CRLR).
15 In the case of RAMP3, it has been shown that together with CRLR it generates an ADM receptor. Upregulation of this gene might play a role in regulating the responsiveness of AtT-20 cells to ADM after CRH exposure or to other extracellular stimuli as it is not known whether RAMP3 might regulate other G-coupled receptors. Although CCK is secreted by AtT-20 cells, induction of its expression by CRH has not
20 been previously reported (13). Interaction between CCK and CRH has however been intensively studied and demonstrated in panic attacks, depression, anxiety and gastric emptying (14-19). Most of these experiments point towards a role for CRH in mediating the central effects of CCK. Our data indicate that CRH in addition might function as a CCK secretagogue. A very similar situation to that of CCK seems to be
25 case for adrenomedullin as well. It has been demonstrated that ADM is expressed in pituitary gland and affects basal and CRH-stimulated ACTH release in animals, thus suggesting its potential role in regulating the hypothalamus-pituitary-adrenal axis(20-23). Current expression data show that CRH induces ADM and CT. In addition recent findings indicate that circulating adrenomedullin is increased in Cushing's disease, and
30 the pituitary gland may represent the site of the elevated production of ADM (24), suggesting CRH might be inducing ADM.
In conclusion, we have unraveled part of the corticotropin-releasing hormone receptor-1 activated gene network and have identified several novel targets of this signaling cascade. Our findings trigger the need for further experiments to elucidate the function
of these transcriptional responses to CRH stimulation both on a cellular and whole organism level.

1. De Souza EB 1995 Corticotropin-releasing factor receptors: physiology, pharmacology, biochemistry and role in central nervous system and immune disorders. *Psychoneuroendocrinology* 20:789-819
- 5 2. Holsboer F 2001 Stress, hypercortisolism and corticosteroid receptors in depression: implications for therapy. *J Affect Disord* 62:77-91
3. Holsboer F, Gerken A, Stalla GK, Muller OA 1987 Blunted aldosterone and ACTH release after human CRH administration in depressed patients. *Am J Psychiatry* 144:229-231
- 10 4. Holsboer F, Gerken A, von Bardeleben U, Grimm W, Beyer H, Muller OA, Stalla GK 1986 Human corticotropin-releasing hormone in depression--correlation with thyrotropin secretion following thyrotropin-releasing hormone. *Biol Psychiatry* 21:601-611
5. Nemeroff CB, Owens MJ, Bissette G, Andorn AC, Stanley M 1988 Reduced corticotropin releasing factor binding sites in the frontal cortex of suicide victims. *Arch Gen Psychiatry* 45:577-579
- 15 6. Raadsheer FC, Hoogendijk WJ, Stam FC, Tilders FJ, Swaab DF 1994 Increased numbers of corticotropin-releasing hormone expressing neurons in the hypothalamic paraventricular nucleus of depressed patients. *Neuroendocrinology* 60:436-444
7. Wouters L, Göhlmann HW, Bijlens L, Kass SU, Molenberghs G, Lewi PJ 2002 Graphical exploration of gene expression data: a comparative study of three multivariate methods. *Biometrics*
- 20 8. Boutillier AL, Monnier D, Lorang D, Lundblad JR, Roberts JL, Loeffler JP 1995 Corticotropin-releasing hormone stimulates proopiomelanocortin transcription by cFos-dependent and -independent pathways: characterization of an AP1 site in exon 1. *Mol Endocrinol* 9:745-755
9. Boutillier AL, Sassone-Corsi P, Loeffler JP 1991 The protooncogene c-fos is induced by corticotropin-releasing factor and stimulates proopiomelanocortin gene transcription in pituitary cells. *Mol Endocrinol* 5:1301-1310
- 25 10. Mazzucchelli C, Sassone-Corsi P 1999 The inducible cyclic adenosine monophosphate early repressor (ICER) in the pituitary intermediate lobe: role in the stress response. *Mol Cell Endocrinol* 155:101-113
11. Oliveira-Dos-Santos AJ, Matsumoto G, Snow BE, Bai D, Houston FP, Whishaw IQ, Mariathasan S, Sasaki T, Wakeham A, Ohashi PS, Roder JC, Barnes CA, Siderovski DP, Penninger JM 2000

Regulation of T cell activation, anxiety, and male aggression by RGS2. *Proc Natl Acad Sci U S A* 97:12272-12277

12. Tse A, Lee AK 2000 Voltage-gated Ca²⁺ channels and intracellular Ca²⁺ release regulate exocytosis in identified rat corticotrophs. *J Physiol* 528 Pt 1:79-90.:79-90
- 5 13. Beinfeld MC 1992 CCK mRNA expression, pro-CCK processing, and regulated secretion of immunoreactive CCK peptides by rat insulinoma (RIN 5F) and mouse pituitary tumor (AtT-20) cells in culture. *Neuropeptides* 22:213-217
14. Coskun T, Bozkurt A, Alican I, Ozkutlu U, Kurtel H, Yegen BC 1997 Pathways mediating CRF-induced inhibition of gastric emptying in rats. *Regul Pept* 69:113-120
- 10 15. Kellner M, Wiedemann K, Yassouridis A, Levengood R, Guo LS, Holaboer F, Yehuda R 2000 Behavioral and endocrine response to cholecystokinin tetrapeptide in patients with posttraumatic stress disorder. *Biol Psychiatry* 47:107-111
- 15 16. Geraciotti TD, Jr., Ekhtor NN, Nicholson WE, Arndt S, Loosen PT, Orth DN 1999 Intra- and inter-individual correlations between cholecystokinin and corticotropin-releasing hormone concentrations in human cerebrospinal fluid. *Depress Anxiety* 10:77-80
17. Calogero AE, Nicolesi AM, Moncada ML, Coniglione F, Vicari E, Polosa P, D'Agata R 1993 Effects of cholecystokinin octapeptide on the hypothalamic-pituitary-adrenal axis function and on vasopressin, prolactin and growth hormone release in humans. *Neuroendocrinology* 58:71-76
- 20 18. Biro E, Samyai Z, Penke B, Szabo G, Telegdy G 1993 Role of endogenous corticotropin-releasing factor in mediation of neuroendocrine and behavioral responses to cholecystokinin octapeptide sulfate ester in rats. *Neuroendocrinology* 57:340-345
19. Kamilaris TC, Johnson EO, Calogero AE, Kalogeras KT, Bernardini R, Chrousos GP, Gold PW 1992 Cholecystokinin-octapeptide stimulates hypothalamic-pituitary-adrenal function in rats: role of corticotropin-releasing hormone. *Endocrinology* 130:1764-1774
- 25 20. Shan J, Krukoff TL 2001 Intracerebroventricular adrenomedullin stimulates the hypothalamic-pituitary-adrenal axis, the sympathetic nervous system and production of hypothalamic nitric oxide. *J Neuroendocrinol* 13:975-984
21. Martinez V, Cutitta F, Tache Y 1997 Central action of adrenomedullin to inhibit gastric emptying in rats. *Endocrinology* 138:3749-3755

22. Parkes DG, May CN 1995 ACTH-suppressive and vasodilator actions of adrenomedullin in conscious sheep. *J Neuroendocrinol* 7:923-929
23. Samson WK, Murphy T, Schell DA 1995 A novel vasoactive peptide, adrenomedullin, inhibits pituitary adrenocorticotropin release. *Endocrinology* 136:2349-2352
- 5 24. Letizia C, Di Iorio R, De Toma G, Marinoni E, Cerci S, Celli M, Subioli S, D'Erasmo E 2000 Circulating adrenomedullin is increased in patients with corticotropin-dependent Cushing's syndrome due to pituitary adenoma. *Metabolism* 49:760-763

SEQUENCE LISTING

SEQ ID NO.1

5 atgcagctgagaaaaatgcagaccatcaaaaaggagcccgacccctagatcctaccagcagctcagacaagatgctgctgctgaactctgcttagct
 gaggtgcccagggacctagcctcaggtgaagattgctcctgaacgaaggagcattggggaanaaanaatcctcggcgtgctggaganaaacgggaat
 tcattccggacgagaaanaagacgccatgtattgggagaaacggcgganaaanaacgaagctgccaaaagatctcgggagaaagcggcgcctcaatg
 acctgggtttgggagaaagcgtgattgccctgggagagaaaatgccacttanaaagctgagctgctcctcctgaaattaaagtgttgaattagctccac
 ggctatgcccagaanaatccagaanaacagtaattccacagctgcttacttccagactaccagacatccaanaggctgccgtgagctctttgtggacgagc
 10 atgagcctcgcgattgtagccgggaagttgcatctcagtcacaaagcncctctccccagagctcgcctccgaltgtctagagggtgtctcgggtgagcacac
 tcaggaagagccccgcacaggggggctgcccggagccctgagaaanaagttccctgtgatcaagcaggagcccgtggaggttggagagccttggcaggga
 ggccaaggagaggagcgggggacgtatccacctccatcaccagagctacatgggaagcttttccacttactccactccccacccctcttgcaggctc
 catgggtccactagcaactcccccaagaacctcagaggccgatgaggggtgtagtgggcaagcttctgaltggggaagacgaacaacaggttccctaagg
 gcccacatccattctccagtgagctgcaaacgggttcacgccacgggtgtaggggttccggaagtgaaccttctgcttaccgcacaagcttccgaltta
 15 agccaaaggccatgcaggtgcaanaaggaggttggagacagggaggttgaaggcatgcagaaactcttccacccggcgtgacgacccaanaagacatt
 tgacctggagaaacatggaacctcgggtatggccattcctccctcccttcttccagtgaggtgacgaacatcaagatttggctcccaaatcggaac
 actggcatcacaaagaactgagcagcaanaactcagagtagcttcaaaacaggtgtgtgggaaggtcaaaagacgggtgctataaggttccgaagcttgag
 aattgtatttgaagcaggggaatagcaacttatctgcagaggtgtgtcgtcnaagagattcatagccacacacacccgatctcggcttccgactccaggtaa

20 SEQ ID No.2

25 MQLRKMQTIKKEPAPLDPTSSSDKMLLLNSALAEVAEDLASGBDLLNEGSMGKNKSSACRRK
 REFIPDEKDDAMYWBKRRKNNEAAKRSREKRRRLNDLVLENKLIALGEENATLKAELLSLKIKFG
 LISSTAYAQRIQKLSNSTAVYFQDYQTSKAAVSSFVDEHEPAMVAGSCISVVIKHSFQSSLSVDSEV
 SSVEHTQESPAQGGCRSPENKFPVIKQEPVELESFAREAREERGTYSTSYQSYMGSFSSTYSHSPF
 LLQVHGSTSNSPRTSEADEGVVGKSSDGEDEQVFKGPIHSPVELQRVHATVVKVPEVNPSPALP
 HKLRIKAKAMQVKVEALDSHFEGMKLSSPADIAKRHFDLBKHOTSQMAHSSLPFVSQVTNI
 QDWSLKSEHWHKELSSKTQSSFKTGVVEVKDGGYKVSEANLYLKQGLANLSAEVVSILKRFT
 30 ATPISASDSR

30 SEQ ID No. 3

35 tgtccgctctgcttccacacctagcaccacccagcccgctgctgccccgggtgagaacccccagcttgggcttgcattggtccagcagggtggccctgag
 ctcttgacagggggcttgcctatagacctgcagggccctgagggcctcagactacacactcaaggggcaaggagggccctggtggccacctaagagccacctct
 gtcccccggccctgctgccccactgactgctgactgagaccagcagtgacccctgagctgctgccccactgctcctcctggtccctgaggttggtctgtc
 cgaggacggagcactctctgaagcagcggcctaacgggaagcagcccccaaggccctccaccgcagcatgggcagtgcccagccagggcctgagcaacg
 tgcctccgggttgcctgctactgttccagatgtggcaccacgaacaggagcggagagcagcatcaggagcaatgggcccctgagaaagcagggaatg
 40 ggtctctagctccaccggccacccctgagcagggccctgtctgcttttactctcttaatttaacatgtatcccgacgatagcagcttgggtcgcacaagtcc
 cgaggcccggtgcccggggagggaccacccggaggaagcccggagcaggtgtcccgtcattgacagccaggccctctgggagcaggttggatgagcacttgc
 agagcaggtgcaatcgtatgttggggcgagggtcctgaaagatattgagacggcctgcaagcttctgaacatcacagcagaccctgggggactggagcc
 ctggtaacgtgcagaaaggccttatggacagaacaccagtaaccggctgcccacagcaggaaggccctccagggagctggcggttaaggagctgtgc
 ggcattgtccgaggaaacagttccgtcagcgtgcacccctgggtgggagttactgcattgcccacctgggacatctgggaagctggccttgatgaaggga
 45 gaggacctcgcctgggaccccttactactgcgcctccaccagcgaaggagggctggagggatgttgaggtggagctgctgctccgggagccatt
 caccgtggcagttcctgaagaaactgctgctcnaagccccacagctaltggccgcttactccgtggctcaacaaaggagaanggcattctcaaaattgagg
 actcagcaagggtggcccgaactgtgggggtgtgcgcaagaacgggccaagccatgaactatgataaactaagccgctccatccggcattatcaagaa
 ggcatcattcgtaaacccgacatctctcagcggctgtctaccaattgtgcatccagctgagagccacagagaccagggcctacacactgccccagg
 50 cagccactctctgttggcctggctctctgctcactctgaattcaggggctgctgcttntccagaacccaaagggtccagatagacagccactgacttag
 ggatacacatgagctctctgggtcatacacaggccccaggaagatcagggggcctggttcagcacacaggggacttgaccaagtcagctcaccgggaca
 gtagatgactgctgctgctcctgcccacaaatccgtgacatattcggcatggtgtcaagagatgtctgacctgctggttgggaaggcagggtgcccgtggg
 gatggataataagacgtatgataactg

SEQ ID No.4

55 MGSASPGLSNVSPGCLLLFPDVPARTGTBKAASGAMGPEKQEWSPSPATPEQGLSAFYLSYFN
 MYPDDSSWVAKVPEARAGEDHPEEPEQCPVIDSQASGSLDEHSLEQVQSMVVGEVLKDIETAC
 KLLNITADPGDWSFGNVQKWLWTEHQYRLPPAGKAFQELGGKELCAMSEEQFRQRAPLGGD

[illegible]

SEQ ID No.6

5

PRDKLT LGKPLGEGCFGQVVM AEAVGIDKDKPKEAVTVAVKMLKDDATEKDLSDLVSEMEM
MKMIGKHKNINLLGACTQDGPLYVIVEYASKGNLREYLRARRPFGMEYSYDINRVPEEQMTFK
DLVSCYQLARGMEYLASQKCUHRDLAARNVLVTENNVMKIADFLARDINNIDYYKKT TNOR
LPVKWMAPEALFDRVYTHQSDVWSFGVLMWEIFTLGGSPYPOIPVEELFKLLKEGHRMDKPTN
CTNELYMMMRDCWHA VPSQRFTFKQLVEDLDRILTLTNEEYLDLTQPLEQYSPSYPTDSSSCS
SGDSDVSFDPMPFYEPCLPQYPHINGSVKT

[illegible]

45 MTVKAEEAARSTLTYSRMRGMVAIIAFMKQRRMGLNDFIQKIASNTYACKHAEVQSILKMSHP
QPELMNANPSPPPSPSQQINLGPSSNPHAKPSDFHFLKVGKGSFGKVLARHKAAEEVFYAVKV
LQKKAILKKKEEKHIMSERNVLLKNVKHPFLVGLHFSFQTADKLYFVLDDYINGGELFYHLQER
CFLEPRARFYAABIASALGYLHSLNIVYRDLKPENILDSQGHVLTDFGLCKENIEHNGTTSTFCG
TPEYLAPEVLHKQPYDRTVDWWCLGAVLYBMLYGLPPFYSRNTAEMYDNLNKPLQLKPNITN
50 SARHLEGLLQKDRTKRLGAKDDFMEEKSHIFFSYINWDDLINKKITPPFNPNVSGPSDLRHFDPE
FTEPSPSPSIGRSPDSILVTASGKDEAAEAFLGFSYAPPVDSFL

55 gcaaggcggcggtgagagcgcgcgtgaaagccgcgcggaacgcgcgtgcacctccgcgactctactacggcgaagctagtctcggacggggctgctgctcccgcgcgcacaccagccctctgggtgaaacgacagggagcgtccggcctccccagcaccggccctgcgagactcaaaaacagccacaccgcacaaaggcgaagcctcgggctggaaaggagcgcgagccttcagcgcgcgcgcctccgcgggaaggatcacatctccgtgtctcaaaaaccccggggcgagagcgcgcgcggggcgcgtg

[illegible]

SEO ID No. 14

25 MLLSKFGSLAHLCPGGVDHLPVKILQPAKADKESFEKVYQVGAVLGSOGFGTVYAGSRIADGL
PVAVKHVVKERVTEWGS LGGVA VPLEVLLRKVGAAGGARGVIRLLDWFERPDGFLVLERPE
PAQDLDFITERGALDEPLARFFAQVLA AVRHCINCGVVHROIKDENLLVDLRSGELKLDGFS
30 GAVLKDTVYTDFDGRVYSPPEWIRYHRYHGRSATVWSLGVLLYDMVCGDIPFEQDEELRGR
LFFRRRVSPQCQLIEWCLSLRPSERPSLDQIAAHPWMLOTEGSVPENCDLRLCALD TDDGASTT
SSFRSL

SEQ ID No.15

[illegible]

5

10

15

20

25

30

3.

4

4

SEQ ID No.18

5

3

MELLRTITYQPAAGTKMCEQALGKACGGDSKKKRPQQPSGDPQPAQVTPAAPHHHHHS
GPEISRIIVDPTTGKRYCRGKVLGKGGFACKEYMTDLTNNKVYAANKIPIHSRAKPHQREKIDKE
IELHRLHLHKHVVFYHYFEDKENIYILLEYSRRSMAHILKARKVLTEPEVRYYLRLQIVSGLKY
LHEQEILHRDLKLGNFINEAMBLKVGDFGLAARLEPLEHRRRTICGTPNYLSPEVLNKGHGCE
SDIWALGCVMYTMLLGRPPFETTNLKETYRCIREARYTMPSSLLAPAKHLIASMLSKNPEDRPSL
DDIIRHDFELQGFTPDRLSSSCCHTVPDFELSSPAKNFFKAAALFGGKKDKARYNDTHINKVSK
EDEDIYKLRHDLKKVSITQPSKHRADDEEPQPPPTTVARSQTSAVENKQQIGDAIRMTVRGTLGS
CSSSSECLEDSMTGMSVADTVARVLRCLENNPEADCIPEQLSTSFEQVWTKWVDYSNKYGFY
QLSDHTYGVLFNNGAHMSLLPDKKTVHYIYABLGCQSVFPATDAPEQFISQVTVLKYFESHYMBE
NLMDGGDLPSVTDIRRPRLYLLQWLKSDKALMMLFNDGTFQVNFYHDHTKHCNQSEEYLLTYI
NEDRISTTFRLTTLMSGCSLELKNRMEYALNMLLQRCN

SEQ ID No. 25

5 aaaggcagcctgataaagctccnctgacagggctgtctgcccagctccagctatgctcctcttgcctctgaaagtgcctcaggatgaaaccacagctccca
aattagcctgggaaagagtgtgcgggaccagcagcctttaaaccgctgagtgcttgcctatgttcaagactgctgtttggatgggtaafgctagctagc
10 actccatcagacatgacagcnaaaatctccaaaagaatttactgcttggaaatctgaggtttgcataaagacttcaaggagcagatgcgcttggaaact
tgagcttccaaagctaccaggaacagacctacatctccaaaattctccacgagcttccaaaggaaattcccatgcttttcagaaagtgtgctggaat
aanaagcatccgacagcggcgtgcttcaoggtgggtctacatgctttgagtggaanaatggcccttccaggtcggagcccatgggaccttcaagcc
ggctcttctgctgggactgggtacttcatgcccgttcttgggcacagccagcgcagggagcttcttctacgacttggacnccgactatgacttgcacca
aaagcgatgtccaggaactcacttccagtgagcaaacggcgatgacctgtgacttcccttggccaggttctggcagcttgcgaagtgaaga
15 acaacttccaccctgtgacgaaccttcatggagcggcgaacaaagggttaccagcggcctagtcaggctccagctcctcagagctcagcctgcaagagga
atcatatcagaaactagcaatggagcgtgggaagaaactagactgtgcttgaaccnctagagaccatccagacotacogctctgtcagcgaatgg
cttcaancaagttaaaaaggatgctgaacgggagctgacacacctctcagagatgagcagatcagggnaaccaggtgtctgagtaatttcaaacacgtt
cttagacaaagcagaacgatgtggaatcccatctccacgcagaaaggcagggaggaagaaagaaagcagcagctatgacccagataagtggaag
gaagaaactgatgcacagctcagcctgaacaacacagcatctcagcttccgggaacacaggaatagagatcatcagccagagcagctgga
20 gacotgaacaaatggggccttaacatcttcaatgtgctgggtactacataatggcccttccagctgcatctgtatgcaatattccaggaagagacctt
ctgaagacgtttaaatctatctgacaccttgttaacctatcatgactttagaagaccattaccattctgactgtgcatatcacaaacagcctgcatgct
gacgtggccagctcaactacgttctcttctacgcccggcacttggatgactttagaagaccattaccattctgactgtgcatatcacaaacagcctgcatgct
atgctgactatcctggagcttccaatcagtttctcatcaatcaaatcttgaactgttctttagatgataatgatgaactgttcttggaaaacatcacttgcgtg
ggatcnaattgtctacaagaggaacactgcgacatcttgaatattccaaagaaagccagacactcaggaatgggtgattgacatgggtgtg
25 caactgataatgtccaaacacatgagccttctggcagacctttaaanaattgtagaaacaaagaggtgacaaagctccggtgttcttccctgacaaat
tactgaccggatcaggttcttgcgaacatggtacactgtgcagacctgagcaacccaccaaagtcttggaaattgatacggcaatgacccatcgtatca
tggagggagttttccagcagggagacaaagagcgggagagggggaattgagattagcccaatgtgtgataagacacagcttctgtggaataatccag
gttgggttcatgactacattgtccatccactgtgggagacctggggcagacctgttcaaacggatgctcaagatattctgatacactagaaatcaacagg
aactgtgatacagagatgatacccccagagcccttccccgccactggatgagagggagcaggggactgccaaggccctgagagaaattcagtttgaactg
30 acccttgaaggaaggatctgagggaccgggaanaaggagggaagggcccaagctatttcagcagacaaagacgcttctgtgattgacccagga
cagggatcttctggaagagactgacatagacattgcaacagaaagacaaagcttctcagatgacacataatctctcctctgtgtggaatgaacttccac
cctgactgagcactgcccgtgagtgtagggctacactacatggccaaaggcctgcacaggaacaaaggccacctggccttccagttacttgaattgga
ggcagaatgccaggccgtgaagcaaatagcagttccatgtctgttctgcttgcctgcaagcttggcggagacccgcagctgtatgtgtagtagggcc
agttccatcaanaagctaaatgcttgaanaacagaggacacaaagctgagagattgctctgcacttgggttgggaagctgtctgacagatgactgaac
35 tcaatacaacttcatataatctaccaccccaacccatgtctgccaacctgtgtgcttttttgaanaatgttgcgccttgaatgcttggtaantatc
tagagtttagtaccacttctacaaactttttagtcttcttgaanaaanaaaaaaanaaaaaaanaaaaaaanaaaaaaanaaaaaa

SEQ ID No.26

35 MTAKNSPKFTASESEVCYKTFKEQMRLELELPKLPGNRPTSPKISPRSSPRNSPCFFRKLLVNKSI
RQRRRTVTAHTCFDVENQPSPGRSPLDPQAGSSGLVLHAAFFGHSQRRESFLYDLSDYDLSPK
AMSRNSSLPSEQHGDDLIVTPFAQVLA SLRSVRNNFILLNLHGAPNKRSPAASQAPVSRVSLQ
ESYQKLAMETLEELDWCILDQLETIQT YRSVSEMASNKFMRMLNRELTHLSEMSRSGNQVSEYIS
40 NTFLDKQNDVEIPSPTQKDREKKKKQQLMTQISGVKKLMHSSSLNNTSISRFGINTEDEDHLAKE
LEDLNKWLGNIFNVAGYSHNRPLTCIMYAIQFLEILAAJFAAAIHDVDHPGVSNQFLINTNSELALMYN
HNSLHAADVAQSTHVLLSTPALDAVFDTLEILAAJFAAAIHDVDHPGVSNQFLINTNSELALMYN
DES VLENHHLAVGFKLLQEBHCDIFQNLTKKQRQTLRKMVIDMVLATDMSKHMSLLADLKT
VETKKVTSSGVLILLDNYTDRIQVLRNMVHCADLSNPTKSLELYRQWTDRIEEDFQQGDKERE
45 RGMESISPMCDKHTASVEKSQVGFIDYIVHFLWETWADLVQPD AQDILDLEDNRNWNWYQSMIPQ
SPSPPLDERSRDCQGLMEKFQFELTLEEDSEGPEKEGEGHSYFSTKTLCVIDPENRDSLEETDID
IATEDKSPIDY

SEQ ID No.27

50 ggccgcgcgggagctgagaatgcaaaagtccatgttcttggctgtccagcagcagctgcgtacccatggacaagagtgcaggcgaacggcccaaggt
cgaggagaagcgggaggaanaatgaagcggacactttaaaccattggaagaccgttggagctacttcttgcagaattctctgtccttgggaagcccaa
aactggcaagaaaagcaaacgcaaacctttaaagccttctcctgggaagcgcacgcttggcagaaagcatttgaactgtgctggcagtaaat
gggctggctgcagctcagggcgttttaagtcgngtctgtggaagaaacattgaattcgttgggtgtggaagacttcaaaaaacaaatccccaa
aaactgtcttcaaaaagcaggaatctatacogacttatagagaaggaagctccc aaagagataaactagacttcaaacgaaatcttgaattgccc
55 aaaaataccaagaggctacaaagtggctgttccacagctcagaaagaggtgtacagtttgaaggagaacattcttactcctggttcttggatccgaat
tctaccaggacttatgtaaaaagccacagatcaccacggagcccatgtctacatgagaccagagctccccacacacaaaaggacattcattctgtct
ccaagagcaaaaggctgtgacctgccagaaaanaaaaaaanaactgacctgaattcagcctgagtgtaggaataacatgctcagaactattgattcaatg

5

MQSAMFLAVQHDCVPMDKSAGNQPKVEEKREKMKRTLNLHWKTRL SYFLQNSSAPGPKPTGK
KSKQQTFIKPSPEEAHVWABAFDELLASKYGLAAFRFLKSEFCEENIEFWLACEDFKKTKSPQK
LSSKARKIYTD FIEKEAPKEINIDFQTKSLIAQNIQEATSGCFTTAQKR VYSLMENN SYPRFLESEF
YODLCKKPQITTEPHAT

10

13

20

24

30

3.

4

MTMETVESQQDRSVTRSVAEHSSAHMQTGQISVPTLAQVATIAETDDSDADSEVIDSHKRREILSR
RPSYRKILNELSSDVPGIPKIBEEKSEEBGTPPNIATMAVPTSIYQTSTGQYNEETDLAPSHMAAAT
GDMPTYQIRAPFTTALPQGVVMAASPQSLHSPQQLAEEATRKRELRLMKNREAARECRKKKEY
VKCLENRVAVLENQNKTLIEELKALKDLYCHKAE

44

50

[illegible]

5.

[illegible]

SEQ ID No.32

35 MSDKMSSFLHIGDICSLYAEGSTNGFISTLGLVDDRCVVQPEAGDLNPPKKFRDCLFKLCPMNR
YSAQKQFWKAAKPGANSTTDAVLLNKLHHAADLEKKQNETENRKLGLTVIQYGNVQLLHLKS
NKYLTVNKRPLALLEKANMRVTLDEAGNEGSWFYIQPFYKLRSDSVIGDKVVLNPNVAGQ
40 PLHASSHQVLDPNPGCNEVNSVNCNTSWKIVLFMKWSDNKDDILKGGDVVRLFHAEQEKFLTCD
EHRKKQHVFLRTTIGROSATSATSSKALWEVEVVQHDPCRGGAGYWNLSLFRFKHLATGHYLA
EVDPDFEBECLFQPSVDPDQDASRSRLRNAQEKMVYSLVSVPFONDISSIFELDPITLRGGDSL
PRNSYVRLRHLCTNTWVHSTNIPIDKBEEKPVMLKIGTSPLKEDKEAFAIVPVSPAENVRLDFAN
DASKVLGSIAGKLEKGTITQNERRSVTKILLEDLVYFVTGGTNSGQDVLEVVFSPKPNRERQKLM
EQNILKQIFKILLQAPFTDCGDGPMLRLEELGDQRHAPFRHICRLCYRVLHRSQDQYRKNQEBYIAK
45 QFGFMQKQIGYDVLAEDTITALLHNNRKLLEKHITAEDTFVSLVRKNREPRFLDYLSDLCSVM
NKSIPVTQELICKAVLNFTNADILBETKLVLSRFEFEGVSTGENALBAGEDEEVWLFWRDSNKEI
RSKSVRELAQDAKEGQKEDRDILSYRYQLNLFARMCLDRQYLAINESGQLDVLILRCMSDE
NLPYDLRASFCRLMLHMKHVDRLDPQEQVTPVKYARLWSEIPSEIAIDDYDSSGTSKDEIKERFAQT
MEFVBEYLRLDVVCQRFPFSDKEKNKLTFEVVNLARNLIYFGFYNFSDLLRLTKILLAILDCVHVT
50 TIFPISKMTKGEENKGSNVMRSHGVGELMTQVVLRGGGFLPMTFMAAAPEGNVKAEBPEKEDI
MVMDTKLKMEILQFILNVRLDYRISCLLCIFKREFDESNSQSSETSSGNSSQEGPSNVPGALDFEHI
EEQAEGIFGGSEENTPLDLDDHGGRTFLRVLLHLLTMHDYPPVLVSGALQLLFRHFSQRQEVLFQAFK
QVQLLVTSDVDNYKQIKQDLQDLRSIVEKSELVWYKGGQGPDEPMDGASGENEHKKEEGTSK
PLKHESTSSYNRYRVVKEILIRLSKLCVQESASVRKSRLKQQRLLRNMGAAHVLELLQIPYEKAE
55 VTQMQEIMRLAHEFLQNFCAAGQONQALHKKHNLFLKPGILEAVTMQHIFMNNFQLCSEINER
VVQHFVHCIEFHGRNVQYIKFLQTVKAEGKFIKKCQDMVMAELVNSGEDVLVfyndRASfQTL
IOMMRSERDRMDENSPLMYHIHLVELLAVCTEGKNVYTEIKCNSLLPLDDIVRVVTHEDCIPEV

20

SEQ ID No.33

5:

5

20

25

45

50

55

aiaaccnccaaatgaagcaccacaagacaaactgtgttgagcgcccccctggggccctgggcctatcttaatacatattgccatcttactgctctgattcctaggaacc
 ccgcaatgacactgcaataatgtccaccatgcgcacagggcaactgtgtggcatgcacagccacagcagcctgtggaagtatgcctgctgatatgccagsgcaictgatgtgt
 ccagcaagggatccccaaccctgcacaaactccgaaacccgcacactctactgtcttcgggaagagacactgccggcggagcgtgtgtgtccgactccacagactcgg
 ctacactctctctgagctctggaaacacctaactacacagtaagtgtcttaataggggagagcaagggagtgggcgcaagtccaccctggccaacatctttgcaggtgtgtga
 tgaacagcatgtagacagcgactgtgagagctctgggagagagatatacatatgagcgtaccctgctgtgtgacgggaagagagtgcnaaccattatctctctggacatg

5

20

25

30

35

45

50

55

SEQ ID No.47

atggctgaacaacilcttccacaggcttgatitgagcaatatgcggaaagctgtgaagatncgagagagnaccccaagaagacattttcaacctaccaat
 5 gggaicacatctatcatttaaacacatgcaccgatacacgctggagntgticagaacatgccagttitgccacagttccgaggaatcatccacaagacatt
 altgacagaaggtgtccaggcttccctggaaagccagaaagcicacactgggtgtcgtgaagtcaggaaagctcgtggctctgaaaaccaatgggtgatgga
 aactgctcatgcatgacgctgticagatcatgtgggtgttcagagatactgacctggctcgtgaggaaagccctctgagcacccttaaggagacagaca
 ctcggaaactttaaatccgctggcagctgggaatctctgaatactcaggaattgtgaaacaggactttgtacgacactcggaaactgggaatgacgaatgg
 gacaacttgggtcaaaatggcatcagcagacacacctgcagccggaaagtggacttcagatcaattccctgggaagaaatccacatatgttccctcagcaacat
 10 cctcagaagagcccaicattgtcatticagacaaaatgctaagaaagtggaaatcgtgtccaatgttgcctcttgaaagtgggtggattatctgctcttcac
 tggcctgcccaaggaggtgttacagataccatcgtcctaggatgacagccagcactttgtacccctgggtgacccctgaaggacagtggaacctgaacttcg
 cgtgttccacttgaacagagaccgggttgggttgaagacttaaaagtgtcacttctgacagatcctgagaatgagatgaaaggaaaagcttctaaaggga
 gtiacttgatagtggagatccctgtgcaaggctgggacacccgacgactcactgacacacgctgcaaaatggatgaagcttaacttaccnaagaa
 aaaaattggtagacgattactttgagctgttcagcacgaaatacaagaaatggcaggaagacagcgaacaggccaggagagcggcagcatgcgagaa
 ccccttggagccttccacacccacgctatcactcatgataataatgtgaacaccccaactgtccttcttctatgtccgtaacactcagcccttatgcccac
 15 gaatgtcagagagggcgccaaaagaatcagagcnaagctcccaagctgaactcgaagctaggccctgaaggactcccaaggcgtgggacttggctcct
 caaactggagggccgaggaacccgctggagggacattcagccccacccacagaccccagccctttctctcagtgagacacttgaatgaaggtgca
 ggagctcctgggtgccccttttactttgaatgtgcagcatmatggtatctgtgagcgttgccacgcccggcagaaatggccagccacacccgagacccctggga
 aagtgccaaagcctgcccitcaggaatgtcactcggacatttaattggcatctgagatcctgtttcaaaaggactacagcagagccagctccagccctacttc
 20 cagataccctgctcctgtccacaaagctccaaagtctgacccctcacaaactctccaaagctcactccacacttggccaccggactggaatgttcttcc
 ttctggctgctctcccaaggctgcacggactcaggaagacagagcagcaggaagcaagtgcaaggaaagctggctgcatgttgggacttcaganaa
 acaagggttttgcactctatgtttatcgaatacagagaaaaaagcngtctgttactgctctgcgaagagctgggttcccccggccccagggttcaganaa
 atgtccctgctcctgggagggaggtgcggcacactcggagacacatgtttgaagggtactgtcagaaagtgtttatcgaagctcagaaccagagatcc
 atgaagcaggaagacgggaagaaagcgtgagatcaagccagcatagagacatgctcgaactacacaggtagccctcaaggtgaaatgtgcccggg
 25 cctcctgcaagaaacttctggcctgtcagctgaaggaaactctgtatgaggtgacagcacttngccaaacagagtaggttctgtggccaccggggtgagc
 ccacgctgaagagccccctaacacagcgtgcccggggccctgctgtgacacttggcactgccaagtgtatgttactgcaatgagtgctaccagttc
 aagcagatgtatgctaa

SEQ ID No.48

30 MAEQLLPQALYLSNMRKAVKIRERTPEDIFKPTNGIYHFKTMRHYTLEMFRTQCFCPQFREIHK
 ALIDRSVQASLESQKKLNWCREVRLVALKTNQDGNCLMHAACQYMWGVQDIDLVLKALC
 STLKETDTRNFKFRWQLESLSQEFVETGLCYDTRNWNDEWDNLVKMASADTPAARSLQYN
 SLBEIHIFVLSNILLRPIVISDKMLRSLSGSNFAPLKVGGIYLPHPWAQECYRYPVILGYDSQHF
 35 VPLVTLKDSGPFLRAVPLVNRDRGRFEDLKVFHFLDTPENEMKEKLLKEYLIVMEIPVQGWHDGT
 THLINAAKLDEANLPKEINLVDDYFELVQHEYKWKQENSQARRAAHAQNPLEPSTPQLSLMDI
 KCETPNCPFFMSVNTQPLCHECSERRQKNQSKLPKLNLSKLOPEGLPGVGLGSSNWSPEETAGGP
 HSAPPTAPSLFLFSETTAMKCRSPGCPFTLVQHNQFCERCHARQINASHTADPGKCQACLDQVT
 RTFNGICSTCFKRTTAEPSSSLTSSIPASCHQRSKSDPSQLIQSLTPHSCHRTGNVSPSGCLSQAART
 40 PGDRAGTSKCRKACCMYFGTPENKQFCTLCHIEYRENKQSVTASAKAGSPAPRFQNNVFLGRE
 CGTLGSTMHEGYCQKCFIEAQNQRFHEARRTEEQLRSSQHRDMPRTTQVASRLKCARASCKNIL
 ACRLSEELCMECQHLSQRVGSVAHRGEPTPEEPPKQRCRAPACDHFNGAKNGYCNCEYQFKQM
 YG

SEQ ID No. 49

45 ttttttctacttgaataatttttaaaaaaaagagaataatutttatcccttgacaaaataaaagatctataataaatgtttctagaaatataatga
 aaagataatattacaataatcaaatcaatcactgacagcaaaagtggaatgattctcaagaaaggtgaggaaggaatgcttccgggiccgca
 gcaatgtctctggagaggccctcctgtccttcttctccttcaatgaggtgtgtctcttatttaagaaaacctgatacagcagaictaatcagtttaggaagct
 50 ggtattatttgcacgcnaaataatuttttaaaaaaaatctatcaagatccittaaatacaagttcccaatgcacttagaatacagttacccaattac
 aagctcttgcacttctctgtgtgtagctotacogcanggcgtgaggtattgtcgaagtgaagtcgtgctgctcgtg

CLAIMS:

- 5 1. An isolated polynucleotide encoding a protein, which modulates corticotropin releasing hormone (CRH) signaling said protein having an amino acid sequence selected from the group consisting of SEQ ID NO. 52, SEQ ID NO.: 54 and SEQ ID NO.: 56.
- 10 2. An isolated polynucleotide according to claim 1 wherein said polynucleotide is mRNA, DNA or cDNA.
- 15 3. An isolated polynucleotide encoding a protein, which modulates corticotropin releasing hormone (CRH) signaling said polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO. 51, SEQ ID NO.: 53 and SEQ ID NO.: 55.
- 20 4. An isolated polynucleotide encoding a protein, which modulates corticotropin releasing hormone (CRH) signaling said polynucleotide consisting of a nucleic acid sequence selected from the group consisting of SEQ ID NO. 51, SEQ ID NO.: 53 and SEQ ID NO.: 55.
- 25 5. An isolated polypeptide encoding a protein, which modulates corticotropin releasing hormone (CRH) signaling said protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO. 52, SEQ ID NO.: 54 and SEQ ID NO.: 56.
- 30 6. A vector comprising an isolated polynucleotide according to any one of claims 1 to 4.
7. A vector according to claim 6 wherein the polynucleotide is operably linked to an expression control sequence.

8. A host cell capable of expressing a protein, which modulates corticotropin releasing hormone (CRH) signaling, said protein having an amino acid sequence selected from the group consisting of SEQ ID NO. 52, SEQ ID NO.: 54 and SEQ ID NO.: 56.
- 5 9. A host cell according to claim 8 transfected with a vector comprising a regulatory sequence.
- 10 10. A host cell according to claim 8 transfected with a vector according to claims 6 or 7.
11. A method for identifying a compound capable to alter the CRH signalling response in a cell, said method comprising:
 - 15 a) contacting said cell with CRH in the presence and absence of said compound;
 - b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said cell; and
 - c) compare the amount of said protein in the presence and absence of said compound;whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.2, SEQ ID 20 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID 25 NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.
12. A method according to claim 11 wherein the cell is a eukaryotic cell such as the murine pituitary corticotroph-derived adenoma cell line AtT-20.
- 30 13. A method according to claims 11 or 12 wherein the amount of protein that modulates CRH signaling is being determined using an antibody which binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID

NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.

5

14. A method according to claims 11 or 12 wherein the amount of protein that modulates CRH signaling is being determined by assessing the level of gene transcription of a gene encoding an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.

10

15

15. A method according to claim 14 wherein the level of gene transcription is being assessed using a probe which binds to a polynucleotide encoding an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.

20

25

16. A method according to claims 14 or 15 wherein the level of gene expression is analysed using microarray technology.

30

17. A method for identifying a compound capable to alter the CRH signalling response activity in a cell, said method comprising;

a) contacting a cell which expresses at least one protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ

ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50, with said test compound; and

5 b) compare the CRH response activity of said cell in the presence and absence of said compound.

18. A method according to claim 17 wherein the cell is host cell capable to express
10 at least one protein having an amino acid sequence selected from the group
 consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID
 NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ
 ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28,
 SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID
 NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ
15 ID NO.48 and SEQ ID NO.50.

19. A method according to claim 18 wherein the host cell is transfected with at least
 one vector comprising a regulatory sequence.

20. A method according to claim 18 wherein the host cell is transfected with at least
20 one vector comprising a polynucleotide sequence encoding an amino acid
 sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ
 ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ
 ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24,
25 SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID
 NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ
 ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.

21. A method of diagnosing CRH induced depression in an individual said method
30 comprising;

a) obtaining a biological sample of said individual; and

b) determine the amount of at least one protein that modulates corticotropin
releasing hormone (CRH) signaling in said biological sample;

whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.

- 5
- 10 22. A method according to claim 21 wherein the biological sample is a bodily fluid or tissue sample
- 15 23. A method according to claims 21 or 22 wherein wherein the amount of protein that modulates CRH signaling is being determined using an antibody which binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.
- 20
24. A method according to claims 21 or 22 wherein the amount of protein that modulates CRH signaling is being determined by assessing the level of gene transcription of a gene encoding an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.
- 25
- 30
- 35 25. A method according to claim 24 wherein the level of gene transcription is being assessed using a probe which binds to a polynucleotide encoding an amino acid sequence selected from the group consisting of SEQ ID NO.2, SEQ ID 4, SEQ ID NO.6, SEQ ID NO.8, SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ

ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.26, SEQ ID NO.28, SEQ ID NO.30, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38, SEQ ID NO.40, SEQ ID NO.42, SEQ ID NO.44, SEQ ID NO.46, SEQ ID NO.48 and SEQ ID NO.50.

5

26. A method according to claims 24 or 25 wherein the level of gene expression is analysed using microarray technology.

ABSTRACT

STRESS RELATED GENES

The present invention relates generally to therapy and diagnosis of depression. In particular this invention relates to the polypeptides as well as to the polynucleotides encoding these polypeptides, wherein said polypeptides are shown to play a central role in mediating the endocrine response to corticotropin releasing hormone. These polypeptides and polynucleotides are useful in the diagnosis, treatment and/or prevention of depression.

Table 1

List of proteins reported to modulate CHN signaling									
	Gene symbol	SWISS ID No.	Accession	Protein name	Accession	Accession	Accession	Accession	Accession
	transcription factors								
	102855_at	1	2	transcription factor, interferon 3, regulated					U03148
	104019_at	3	4	prostate specific antigen transcription factor					AB016496
	93618_at	5	6	prostate homeobox (Prosp1)					AF022992
	104155_at	7	8	cellular transcription factor 3					U03110
	Alvasea								
	93090_at	9	10	transcription factor 2					U03152
	97890_at	11	12	transcription factor 2					U03152
	98847_at	13	14	prostate homeobox (Prosp1)					U03152
	100133_at	15	16	FYI protein					U03152
	92310_at	17	18	transcription factor 2					U03152
	transcription factors								
	98055_at	19	20	Chaperonin					U03152
	102789_at	21	22	transcription factor 2					U03152
	92533_at	23	24	cellular					U03152
	calcium signaling								
	103810_at	25	26	phosphodiesterase 4B, cAMP specific					U03152
	97844_at	27	28	regulator of G-protein signaling 2					U03152
	100533_at	29	30	cAMP responsive element modulator					U03152
	calcium signaling								
	94977_at	31	32	phosphodiesterase 4B, cAMP specific					U03152
	98592_at	33	34	phosphodiesterase 4B, cAMP specific					U03152
	phosphodiesterase								
	104622_at	35	36	phosphodiesterase 4B, cAMP specific					U03152
	104588_at	37	38	phosphodiesterase 4B, cAMP specific					U03152
	receptor and channel regulators								
	92534_at	39	40	phosphodiesterase 4B, cAMP specific					U03152
	92388_at	41	42	phosphodiesterase 4B, cAMP specific					U03152
	proteases								
	92820_at	43	44	phosphodiesterase 4B, cAMP specific					U03152
	calcium signaling								
	93874_at	45	46	phosphodiesterase 4B, cAMP specific					U03152
	93874_at	47	48	phosphodiesterase 4B, cAMP specific					U03152
	93874_at	49	50	phosphodiesterase 4B, cAMP specific					U03152

Fig. 1




Fig. 2




Fig.3




Fig. 4




Fig. 4 – continued




Fig.4 -- continued





Fig. 4 - continued

Phosphatases

protein tyrosine phosphatase, receptor-type, II

protein tyrosine phosphatase, non-receptor type, 16



Unknowns

LOC137790 (MAGE-503759)

RIKEN cDNA 180002F13 gene

tumor necrosis factor, alpha-induced protein 3







Fig. 5



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.